

L1 ANSWER 2 OF 10 WPIX COPYRIGHT 2007 THE THOMSON CORP on STN
 ACCESSION NUMBER: 2002-547628 [58] WPIX
 DOC. NO. CPI: C2002-155251 [58]
 TITLE: New imidazolone derivatives are HIV-1 reverse
 transcriptase inhibitors used for treating e.g.
 viral diseases, cancer and bacterial and inflammatory
 diseases
 DERWENT CLASS: B03
 INVENTOR: DYMOCK B W; JONES P S; MERRETT J H; PARKES K E;
 PARKES K E B; PARRATT M J; WALTER D S; DYMOCK W; JONES S;
 MERRETT H; PARKES E; PARRATT J; WALTER S
 PATENT ASSIGNEE: (DYMO-I) DYMOCK B W; (HOFF-C) HOFFMANN LA ROCHE &
 CO AG F; (HOFF-C) HOFFMANN LA ROCHE INC; (JONE-I) JONES
 P S; (MERR-I) MERRETT J H; (PARK-I) PARKES K E; (PARR-
 I) PARRATT M J; (WALT-I) WALTER D S
 COUNTRY COUNT: 96

PATENT INFO ABBR.:

PATENT NO	KIND	DATE	WEEK	LA	PG	MAIN IPC
WO 2002042279	A1	20020530	(200258)*	EN	88[0]	
US 20020107272	A1	20020808	(200259)	EN		
AU 2002021842	A	20020603	(200263)	EN		
US 6518293	B2	20030211	(200314)	EN		
BR 2001015476	A	20030819	(200367)	PT		
EP 1341766	A1	20030910	(200367)	EN		
KR 2003060951	A	20030716	(200381)	KO		
JP 2004514672	W	20040520	(200434)	JA	154	
MX 2003004454	A1	20031001	(200466)	ES		
ZA 2003003690	A	20041027	(200474)	EN	94	
EP 1341766	B1	20051026	(200571)	EN		
CN 1633417	A	20050629	(200574)	ZH		
DE 60114448	E	20051201	(200580)	DE		
ES 2251529	T3	20060501	(200634)	ES		
MX 233501	B	20060106	(200639)	ES		
DE 60114448	T2	20060706	(200645)	DE		
KR 528760	B	20051115	(200682)	KO		

APPLICATION DETAILS:

PATENT NO	KIND	APPLICATION	DATE
WO 2002042279	A1	WO 2001-EP13069	20011112
US 20020107272	A1	US 2001-993848	20011106
US 6518293	B2	US 2001-993848	20011106
BR 2001015476	A	BR 2001-15476	20011112
CN 1633417	A	CN 2001-819352	20011112

DE 60114448 E	DE 2001-614448 20011112
DE 60114448 T2	DE 2001-614448 20011112
EP 1341766 A1	EP 2001-997479 20011112
EP 1341766 B1	EP 2001-997479 20011112
DE 60114448 E	EP 2001-997479 20011112
ES 2251529 T3	EP 2001-997479 20011112
DE 60114448 T2	EP 2001-997479 20011112
BR 2001015476 A	WO 2001-EP13069 20011112
EP 1341766 A1	WO 2001-EP13069 20011112
JP 2004514672 W	WO 2001-EP13069 20011112
MX 2003004454 A1	WO 2001-EP13069 20011112
EP 1341766 B1	WO 2001-EP13069 20011112
DE 60114448 E	WO 2001-EP13069 20011112
MX 233501 B	WO 2001-EP13069 20011112
DE 60114448 T2	WO 2001-EP13069 20011112
AU 2002021842 A	AU 2002-21842 20011112
JP 2004514672 W	JP 2002-544414 20011112
ZA 2003003690 A	ZA 2003-3690 20030513
MX 2003004454 A1	MX 2003-4454 20030520
MX 233501 B	MX 2003-4454 20030520
KR 2003060951 A	KR 2003-706881 20030521
KR 528760 B	WO 2001-EP13069 20011112
KR 528760 B	KR 2003-706881 20030521

FILING DETAILS:

PATENT NO	KIND		PATENT NO	
DE 60114448	E	Based on	EP 1341766	A
ES 2251529	T3	Based on	EP 1341766	A
DE 60114448	T2	Based on	EP 1341766	A
AU 2002021842	A	Based on	WO 2002042279	A
BR 2001015476	A	Based on	WO 2002042279	A
EP 1341766	A1	Based on	WO 2002042279	A
JP 2004514672	W	Based on	WO 2002042279	A
MX 2003004454	A1	Based on	WO 2002042279	A
EP 1341766	B1	Based on	WO 2002042279	A
DE 60114448	E	Based on	WO 2002042279	A
MX 233501	B	Based on	WO 2002042279	A
DE 60114448	T2	Based on	WO 2002042279	A
KR 528760	B	Previous Publ	KR 2003060951	A
KR 528760	B	Based on	WO 2002042279	A

PRIORITY APPLN. INFO: GB 2000-28484 20001122

AN 2002-547628 [58] WPIX

AB WO 2002042279 A1 UPAB: 20060202

NOVELTY - Imidazolone derivatives (I) are new.

DETAILED DESCRIPTION - Imidazolone derivatives of formula (I),

their hydrolyzable esters, ethers and salts, are new.

R1 = H, cycloalkyl, aryl, or alkyl optionally substituted with

optionally substituted phenyl;

R2 = H, alkenyl or alkyl optionally substituted with optionally

substituted phenyl;

R3 = phenyl, benzyl (both optionally substituted), alkyl,

cycloalkyl, or heterocyclyl;
A = alkyl, alkenyl (both optionally substituted),
hydroxyalkyl,
aryl-CH(OH)- or Z-CH₂-Y-CH₂-;
Y = O or NR;
R = H or alkyl;
Z = heterocyclyl or optionally substituted aryl; and
X = S or O;
provided that only one of R₁ and R₂ is H; and when X is O,
then R₁
is not aryl.
INDEPENDENT CLAIMS are also included for:
(1) preparation of a compound of formula (VIII) involving
reacting
a compound of formula (VII) with R₃-XH;
(2) a compound of formula (VIIIa); and
(3) use of (I) for the preparation of a medicament for the
treatment of diseases mediated by the human immunodeficiency virus
(HIV).
R'₃ = phenyl or benzyl (both optionally substituted by 1-5
1-4C
alkyl, 1-4C alkoxy, OH, F, Cl or Br), 7-12C alkyl or 3-8C
cycloalkyl.
ACTIVITY - Anti-HIV; Antibacterial; Antiparasitic;
Virucide;
Anti-HIV; Antiinflammatory; Cytostatic.
In a test to determine anti-HIV antiviral activity (see
Pauwels et
al., 1988, J. Virol Methods 20:309-321), 4-(3,5-dichloro-
phenylsulfonyl)-
isopropyl-1-methyl-5-(pyridin-3-ylmethoxymethyl)-1,3-dihydro-
imidazol-2-
one (Ia) inhibited HIV activity with an IC₅₀ of 5 nM.
MECHANISM OF ACTION - HIV-1-reverse transcriptase enzyme
inhibitor.
In an HIV-1 reverse transcription assay, 4-(3,5-dichloro-
phenylsulfonyl)-3-isopropyl-1-methyl-5-(pyridin-3-
ylmethoxymethyl)-1,3-
dihydro-imidazol-2-one (Ia) inhibited HIV reverse transcriptase
with an
IC₅₀ of 115 nM.
USE - In the preparation of a medicament for the treatment
of
diseases mediated by HIV (claimed); for treating viral diseases
(e.g.
retroviral infection), immune mediated conditions or diseases,
bacterial,
parasitic, inflammatory or hyperproliferative vascular diseases,
tumor and
cancer; and as chemotherapeutic agents, inhibitors of viral
replication
and modulators of the immune system.
ADVANTAGE - The compounds are potent inhibitors of HIV-
reverse
transcriptase enzyme involved in viral replication and have a
potential to
be efficacious as antiviral drugs.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
30 May 2002 (30.05.2002)

PCT

(10) International Publication Number
WO 02/42279 A1

(51) International Patent Classification⁷: **C07D 233/86**,
233/78, 401/12, 401/06, A61K 31/4164, 31/4174, 31/4427

149 Wadnall Way, Knebworth, Hertfordshire SG3 6DT
(GB).

(21) International Application Number: PCT/EP01/13069

(74) Agent: **MEZGER, Wolfgang**; Grenzacherstrasse 124,
CH-4070 Basle (CH).

(22) International Filing Date:
12 November 2001 (12.11.2001)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA,
ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0028484.4 22 November 2000 (22.11.2000) GB

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

(71) Applicant: **F. HOFFMANN-LA ROCHE AG** [CH/CH];
Grenzacherstrasse 124, CH-4070 Basle (CH).


Published:

— with international search report

(72) Inventors: **DYMOCK, Brian, William**; 15 Vesta Av-
enue, St. Albans, Hertfordshire AL1 2PJ (GB). **JONES,**
Philip, Stephen; 58 Digswell rise, Welwyn Garden City,
Hertfordshire AL8 7PW (GB). **MERRETT, John, Her-**
bert; 23 Bush Spring, Baldock, Hertfordshire SG7 6QT
(GB). **PARKES, Kevin, Edward, Burdon**; 60 Hallmead,
Letchworth, Hertfordshire SG6 4BJ (GB). **PARRATT,**
Martin, John; 22 Chapel Lane, Letty Green, Hertford,
Hertfordshire SG14 2PA (GB). **WALTER, Daryl, Simon**;

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: IMIDAZOLONE DERIVATIVES FOR THE TREATMENT OF VIRAL DISEASES



(57) Abstract: The invention describes novel imidazolone derivatives as well as hydrolyz-
able esters or ethers of compounds of formula I and pharmaceutically acceptable salts thereof,
a process for their manufacture, pharmaceutical compositions and the use of such compounds
in medicine (formula I) wherein R¹, R², R³, A and X are as defined in the description. The
compounds of formula I are the human immunodeficiency virus reverse transcriptase enzyme
which is involved in viral replication. Consequently the compounds of this invention may be
advantageously used as therapeutic agents for HIV mediated process.

Imidazolone Derivatives for the Treatment of Viral Diseases

The invention is concerned with novel imidazolone derivatives, a process for their manufacture, pharmaceutical compositions and the use of such compounds in medicine. In particular, the compounds are inhibitors of the human immunodeficiency virus reverse transcriptase enzyme which is involved in viral replication. Consequently the compounds of this invention may be advantageously used as therapeutic agents for the treatment of diseases mediated by the human immunodeficiency virus (HIV).

The disease Acquired Immunodeficiency Syndrome (AIDS) is the end result of infection by the distinct retroviruses, human immunodeficiency virus type-1 (HIV-1) or type-2 (HIV-2). Several critical points in the virus life cycle have been identified as possible targets for therapeutic intervention. Inhibition of one of these, the transcription of viral RNA to viral DNA (reverse transcriptase, RT), has provided a number of the current therapies used in treating AIDS. Inhibition of reverse transcriptase provided the first form of treatment for HIV infection with 3'-azido-3'-deoxythymidine (AZT). Since then several inhibitors have been launched, broadly forming two classes: nucleoside analogues and non-nucleosides. As an example of the latter it has been found that certain benzoxazinones, e.g. efavirenz are useful in the inhibition of HIV RT. However, development of strains of the virus resistant to current RT inhibitors is a constant problem. Therefore, development of compounds effective against resistant strains is an important goal.

Imidazolone derivatives have been described in the literature with different uses in the treatment of the human body.


WO 94/10168 describes imidazolone derivatives and their use as tachykinin receptor antagonists.

WO 96/38421 describes imidazolone derivatives and their use as anti-coagulants.

WO 00/35907 describes imidazolone derivatives and their use in treating benign prostatic hyperplasia.

The object of the present invention is to provide novel compounds which are potent inhibitors of the human immunodeficiency virus reverse transcriptase enzyme, which is involved in viral replication, and which accordingly show a potential to be efficacious as antiviral drugs.

5 This object can be achieved with compounds of formula I



wherein

R¹ is hydrogen, alkyl, cycloalkyl, aryl or alkyl substituted with optionally substituted phenyl;

10 R² is hydrogen, alkenyl, alkyl or alkyl substituted with optionally substituted phenyl;

R³ is alkyl, cycloalkyl, optionally substituted phenyl, optionally substituted benzyl or heterocyclyl;

A is alkyl, hydroxy-alkyl, alkenyl, substituted alkenyl, aryl-CH(OH)-, substituted alkyl, or

15 A signifies a group of formula Z-CH₂-Y-CH₂-,

wherein Y represents O or NR wherein R is hydrogen or alkyl and

Z represents heterocyclyl or optionally substituted aryl;

X represents S or O;

with the provisos that

- 20 (i) only one of R¹ and R² is hydrogen;
(ii) when X is O, then R¹ cannot be aryl;

as well as hydrolyzable esters or ethers of compounds of formula I and pharmaceutically acceptable salts thereof.

The term "alkyl" as used herein denotes an optionally substituted straight or branched chain hydrocarbon residue containing 1 to 12 carbon atoms, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert.-butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl including their different isomers. Preferably, the term
5 "alkyl" denotes an optionally substituted straight or branched chain hydrocarbon residue containing 1 to 7 carbon atoms.

Suitable substituents for the alkyl chain can be selected from one or more of aryl, heterocyclyl, alkoxy, hydroxy or halogen.

Aryl or heterocyclyl as substituents for the alkyl group can also be substituted with 1,
10 2, 3, 4, 5 or where possible more methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert.-butyl, methoxy, ethoxy, n-propyloxy, iso-propyloxy, n-butyloxy, iso-butyloxy, tert.-butyloxy, hydroxy, fluorine, chlorine, bromine or iodine.

Alkyl in R^1 , R^2 and R^3 is preferably an optionally substituted straight or branched chain hydrocarbon residue containing 1 to 7 carbon atoms. Suitable substituents for the
15 alkyl group are selected from aryl, heterocyclyl, hydroxy or halogen. Preferred alkyl groups in R^1 , R^2 and R^3 are unsubstituted straight or branched chain hydrocarbon residues containing 1 to 7 carbon atoms and more preferred alkyl groups in R^1 , R^2 and R^3 are methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert.-butyl or pentyl. Most preferred alkyl group in R^1 , R^2 and R^3 are methyl, ethyl, propyl, isopropyl, n-butyl, sec-
20 butyl, isobutyl, tert.-butyl.

Alkyl for R^1 is preferably isopropyl.

Alkyl for R^2 is preferably methyl.

Alkyl for the substituent A is as defined above, preferably a straight or branched chain hydrocarbon residue containing 1 to 7 carbon atoms. More preferred alkyl group for
25 the substituent A are methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert.-butyl or pentyl.

The term "alkoxy" as used herein denotes an optionally substituted straight or branched chain alkyl-oxy group wherein the "alkyl" portion is as defined above such as methoxy, ethoxy, n-propyloxy, iso-propyloxy, n-butyloxy, iso-butyloxy, tert.-butyloxy,
30 pentyloxy, hexyloxy, heptyloxy including their different isomers. More preferred alkoxy groups within the invention are methoxy, ethoxy, n-propyloxy, iso-propyloxy, n-butyloxy, iso-butyloxy or tert.-butyloxy.

The term "C₁₋₄-alkyl substituted with C₁₋₄-alkoxy, C₁₋₄-alkyl-carbonyl-oxy- or heterocyclyl-oxy" as used herein denotes a C₁₋₄-alkyl, preferably C₁₋₂-alkyl as defined above which may be substituted with 1-3, preferably 1-2 and more preferably on substituents selected from C₁₋₄-alkoxy (preferred C₁₋₂-alkoxy), C₁₋₄-alkyl-carbonyl-oxy- (preferred C₁₋₂-alkyl-carbonyl-oxy-) and heterocyclyl-oxy. Preferred examples are methoxymethyl, ethoxymethyl, methyl-carbonyl-oxy-, ethyl-carbonyl-oxy-, 4-pyridyl-oxy-methyl, 3-pyridyl-oxy-methyl, 2-pyridyl-oxy-methyl.

The term "C₁₋₄-alkyl substituted with optionally substituted phenyl" as used herein denotes a C₁₋₄-alkyl, preferably C₁₋₂-alkyl as defined above which may be substituted with a phenyl group or with a substituted phenyl group which may be substituted with 1,2,3,4 or 5 substituents, preferably 1, 2 or 3 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine or bromine, preferably methyl, ethyl, methoxy, ethoxy, hydroxy, fluorine, chlorine or bromine. Examples are phenylmethyl (benzyl), phenylethyl, phenylpropyl, phenylbutyl, tolylmethyl, tolylethyl, tolylpropyl, tolylbutyl, 2,3-dimethylphenylmethyl, 2,4-dimethylphenylmethyl, 2,5-dimethylphenylmethyl, 2,6-dimethylphenylmethyl, 3,4-dimethylphenylmethyl, 3,5-dimethylphenylmethyl, 3,6-dimethylphenylmethyl, 2-methoxy-phenylmethyl, 3-methoxy-phenylmethyl, 4-methoxy-phenylmethyl, methoxy-phenylethyl, methoxy-phenylpropyl, methoxy-phenylbutyl, dimethoxy-phenylmethyl, dimethoxy-phenylethyl, dimethoxy-phenylpropyl, dimethoxy-phenylbutyl, 2-hydroxyphenylmethyl, 3-hydroxyphenylmethyl, 4-hydroxyphenylmethyl, 2,3-dihydroxyphenylmethyl, 2,4-dihydroxyphenylmethyl, 2,5-dihydroxyphenylmethyl, 2,6-dihydroxyphenylmethyl, 3,4-dihydroxyphenylmethyl, 3,5-dihydroxyphenylmethyl, 3,6-dihydroxyphenylmethyl, 2-hydroxyphenylethyl, 3-hydroxyphenylethyl, 4-hydroxyphenylethyl, 2-hydroxyphenylpropyl, 3-hydroxyphenylpropyl, 4-hydroxyphenylpropyl, 2-hydroxyphenylbutyl, 3-hydroxyphenylbutyl, 4-hydroxyphenylbutyl, 2-fluorophenylmethyl, 3-fluorophenylmethyl, 4-fluorophenylmethyl, 2,3-difluorophenylmethyl, 2,4-difluorophenylmethyl, 2,5-difluorophenylmethyl, 2,6-difluorophenylmethyl, 3,4-difluorophenylmethyl, 3,5-difluorophenylmethyl, 3,6-difluorophenylmethyl, 2-fluorophenylethyl, 3-fluorophenylethyl or 4-fluorophenylethyl, 2-chlorophenylmethyl, 3-chlorophenylmethyl, 4-chlorophenylmethyl, 2,3-dichlorophenylmethyl, 2,4-dichlorophenylmethyl, 2,5-dichlorophenylmethyl, 2,6-dichlorophenylmethyl, 3,4-dichlorophenylmethyl, 3,5-dichlorophenylmethyl, 3,6-dichlorophenylmethyl, 2-chlorophenylethyl, 3-chlorophenylethyl, 4-chlorophenylethyl, 2-bromophenylmethyl, 3-bromophenylmethyl, 4-bromophenylmethyl, 2,3-dibromophenylmethyl, 2,4-dibromophenylmethyl, 2,5-dibromophenylmethyl, 2,6-dibromophenylmethyl, 3,4-dibromophenylmethyl, 3,5-dibromophenylmethyl, 3,6-dibromophenylmethyl, 2-bromophenylethyl, 3-bromophenylethyl or 4-bromophenylethyl.

In case more than one substituent is attached to the phenyl group, these substituents can be identical or different from each other and are also encompassed by the scope of the present invention. For example dimethoxy-phenylmethyl means that both methoxy substituents may be attached to the phenyl ring in the 2,3-position, the 2,4-position, the 2,5-position, the 2,6-position, the 3,4-position, the 3,5-position and the 3,6-position.

C₁₋₄-alkyl groups substituted with optionally substituted phenyl in R¹ are as defined above and preferably phenylmethyl (benzyl), phenylethyl, tolylmethyl, tolylethyl, 2,3-dimethylphenylmethyl, 2,4-dimethylphenylmethyl, 2,5-dimethylphenylmethyl, 2,6-dimethylphenylmethyl, 3,4-dimethylphenylmethyl, 3,5-dimethylphenylmethyl, 3,6-dimethylphenylmethyl, 2-methoxy-phenylmethyl, 3-methoxy-phenylmethyl, 4-methoxy-phenylmethyl, methoxy-phenylethyl, dimethoxy-phenylmethyl, dimethoxy-phenylethyl, 2-hydroxyphenylmethyl, 3-hydroxyphenylmethyl, 4-hydroxyphenylmethyl, 2,3-dihydroxyphenylmethyl, 2,4-dihydroxyphenylmethyl, 2,5-dihydroxyphenylmethyl, 2,6-dihydroxyphenylmethyl, 3,4-dihydroxyphenylmethyl, 3,5-dihydroxyphenylmethyl, 3,6-dihydroxyphenylmethyl, 2-hydroxyphenylethyl, 3-hydroxyphenylethyl, 4-hydroxyphenylethyl, 2-fluorophenylmethyl, 3-fluorophenylmethyl, 4-fluorophenylmethyl, 2,3-difluorophenylmethyl, 2,4-difluorophenylmethyl, 2,5-difluorophenylmethyl, 2,6-difluorophenylmethyl, 3,4-difluorophenylmethyl, 3,5-difluorophenylmethyl, 3,6-difluorophenylmethyl, 2-fluorophenylethyl, 3-fluorophenylethyl or 4-fluorophenylethyl, 2-chlorophenylmethyl, 3-chlorophenylmethyl, 4-chlorophenylmethyl, 2,3-dichlorophenylmethyl, 2,4-dichlorophenylmethyl, 2,5-dichlorophenylmethyl, 2,6-dichlorophenylmethyl, 3,4-dichlorophenylmethyl, 3,5-dichlorophenylmethyl, 3,6-dichlorophenylmethyl, 2-chlorophenylethyl, 3-chlorophenylethyl, 4-chlorophenylethyl, 2-bromophenylmethyl, 3-bromophenylmethyl, 4-bromophenylmethyl, 2,3-dibromophenylmethyl, 2,4-dibromophenylmethyl, 2,5-dibromophenylmethyl, 2,6-dibromophenylmethyl, 3,4-dibromophenylmethyl, 3,5-dibromophenylmethyl, 3,6-dibromophenylmethyl, 2-bromophenylethyl, 3-bromophenylethyl or 4-bromophenylethyl. Most preferred C₁₋₄-alkyl substituted with optionally substituted phenyl in R¹ are phenylmethyl (benzyl), phenylethyl, tolylmethyl, tolylethyl, 2,3-dimethylphenylmethyl, 2,4-dimethylphenylmethyl, 2,5-dimethylphenylmethyl, 2,6-dimethylphenylmethyl, 3,4-dimethylphenylmethyl, 3,5-dimethylphenylmethyl or 3,6-dimethylphenylmethyl.

C₁₋₄-alkyl groups substituted with optionally substituted phenyl in R² are as defined above and preferably phenylmethyl (benzyl), phenylethyl, tolylmethyl, tolylethyl, 2,3-dimethylphenylmethyl, 2,4-dimethylphenylmethyl, 2,5-dimethylphenylmethyl, 2,6-dimethylphenylmethyl, 3,4-dimethylphenylmethyl, 3,5-dimethylphenylmethyl, 3,6-dimethylphenylmethyl, 2-methoxy-phenylmethyl, 3-methoxy-phenylmethyl, 4-methoxy-

phenylmethyl, methoxy-phenylethyl, dimethoxy-phenylmethyl, dimethoxy-phenylethyl, 2-hydroxyphenylmethyl, 3-hydroxyphenylmethyl, 4-hydroxyphenylmethyl, 2,3-dihydroxyphenylmethyl, 2,4-dihydroxyphenylmethyl, 2,5-dihydroxyphenylmethyl, 2,6-dihydroxyphenylmethyl, 3,4-dihydroxyphenylmethyl, 3,5-dihydroxyphenylmethyl, 3,6-dihydroxyphenylmethyl, 2-hydroxyphenylethyl, 3-hydroxyphenylethyl, 4-hydroxyphenylethyl, 2-fluorophenylmethyl, 3-fluorophenylmethyl, 4-fluorophenylmethyl, 2,3-difluorophenylmethyl, 2,4-difluorophenylmethyl, 2,5-difluorophenylmethyl, 2,6-difluorophenylmethyl, 3,4-difluorophenylmethyl, 3,5-difluorophenylmethyl, 3,6-difluorophenylmethyl, 2-fluorophenylethyl, 3-fluorophenylethyl or 4-fluorophenylethyl, 2-chlorophenylmethyl, 3-chlorophenylmethyl, 4-chlorophenylmethyl, 2,3-dichlorophenylmethyl, 2,4-dichlorophenylmethyl, 2,5-dichlorophenylmethyl, 2,6-dichlorophenylmethyl, 3,4-dichlorophenylmethyl, 3,5-dichlorophenylmethyl, 3,6-dichlorophenylmethyl, 2-chlorophenylethyl, 3-chlorophenylethyl, 4-chlorophenylethyl, 2-bromophenylmethyl, 3-bromophenylmethyl, 4-bromophenylmethyl, 2,3-dibromophenylmethyl, 2,4-dibromophenylmethyl, 2,5-dibromophenylmethyl, 2,6-dibromophenylmethyl, 3,4-dibromophenylmethyl, 3,5-dibromophenylmethyl, 3,6-dibromophenylmethyl, 2-bromophenylethyl, 3-bromophenylethyl or 4-bromophenylethyl. Most preferred C₁₋₄-alkyl substituted with optionally substituted phenyl in R² are phenylmethyl (benzyl), phenylethyl, tolylmethyl, tolylethyl, 2,3-dimethylphenylmethyl, 2,4-dimethylphenylmethyl, 2,5-dimethylphenylmethyl, 2,6-dimethylphenylmethyl, 3,4-dimethylphenylmethyl, 3,5-dimethylphenylmethyl, 3,6-dimethylphenylmethyl, methoxy-phenylmethyl, methoxy-phenylethyl, dimethoxy-phenylmethyl or dimethoxy-phenylethyl.

C₁₋₄-alkyl groups substituted with optionally substituted phenyl for the substituent A are as defined above, preferably phenylmethyl (benzyl), phenylethyl, tolylmethyl, tolylethyl, 2,3-dimethylphenylmethyl, 2,4-dimethylphenylmethyl, 2,5-dimethylphenylmethyl, 2,6-dimethylphenylmethyl, 3,4-dimethylphenylmethyl, 3,5-dimethylphenylmethyl, 3,6-dimethylphenylmethyl, 2-methoxy-phenylmethyl, 3-methoxy-phenylmethyl, 4-methoxy-phenylmethyl, methoxy-phenylethyl, dimethoxy-phenylmethyl, dimethoxy-phenylethyl, 2-hydroxyphenylmethyl, 3-hydroxyphenylmethyl, 4-hydroxyphenylmethyl, 2,3-dihydroxyphenylmethyl, 2,4-dihydroxyphenylmethyl, 2,5-dihydroxyphenylmethyl, 2,6-dihydroxyphenylmethyl, 3,4-dihydroxyphenylmethyl, 3,5-dihydroxyphenylmethyl, 3,6-dihydroxyphenylmethyl, 2-hydroxyphenylethyl, 3-hydroxyphenylethyl, 4-hydroxyphenylethyl, 2-fluorophenylmethyl, 3-fluorophenylmethyl, 4-fluorophenylmethyl, 2,3-difluorophenylmethyl, 2,4-difluorophenylmethyl, 2,5-difluorophenylmethyl, 2,6-difluorophenylmethyl, 3,4-difluorophenylmethyl, 3,5-difluorophenylmethyl, 3,6-difluorophenylmethyl, 2-fluorophenylethyl, 3-fluorophenylethyl or 4-fluorophenylethyl, 2-chlorophenylmethyl, 3-chlorophenylmethyl,

4-chlorophenylmethyl, 2,3-dichlorophenylmethyl, 2,4-dichlorophenylmethyl, 2,5-dichlorophenylmethyl, 2,6-dichlorophenylmethyl, 3,4-dichlorophenylmethyl, 3,5-dichlorophenylmethyl, 3,6-dichlorophenylmethyl, 2-chlorophenylethyl, 3-chlorophenylethyl, 4-chlorophenylethyl, 2-bromophenylmethyl, 3-bromophenylmethyl, 4-bromophenylmethyl, 2,3-dibromophenylmethyl, 2,4-dibromophenylmethyl, 2,5-dibromophenylmethyl, 2,6-dibromophenylmethyl, 3,4-dibromophenylmethyl, 3,5-dibromophenylmethyl, 3,6-dibromophenylmethyl, 2-bromophenylethyl, 3-bromophenylethyl or 4-bromophenylethyl. Most preferred C₁₋₄-alkyl substituted with optionally substituted phenyl for the substituent A are phenylmethyl (benzyl), phenylethyl, tolylmethyl, tolylethyl, 2,3-dimethylphenylmethyl, 2,4-dimethylphenylmethyl, 2,5-dimethylphenylmethyl, 2,6-dimethylphenylmethyl, 3,4-dimethylphenylmethyl, 3,5-dimethylphenylmethyl, 3,6-dimethylphenylmethyl, 2-hydroxyphenylmethyl, 3-hydroxyphenylmethyl, 4-hydroxyphenylmethyl, 2,3-dihydroxyphenylmethyl, 2,4-dihydroxyphenylmethyl, 2,5-dihydroxyphenylmethyl, 2,6-dihydroxyphenylmethyl, 3,4-dihydroxyphenylmethyl, 3,5-dihydroxyphenylmethyl, 3,6-dihydroxyphenylmethyl, 2-hydroxyphenylethyl, 3-hydroxyphenylethyl or 4-hydroxyphenylethyl.

The term "cycloalkyl" as used herein denotes an optionally substituted cycloalkyl group containing 3 to 8 carbon atoms, e.g. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl.

Suitable substituents for cycloalkyl can be selected from those named for alkyl, in addition however an oxo group (=O) can be added to the selection.

Cycloalkyl in R¹ is as defined above and preferably cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, preferably cyclopropyl or cyclobutyl and most preferred cyclopropyl.

Cycloalkyl in R³ is as defined above, preferably cyclopentyl, cyclohexyl or cycloheptyl and most preferred cyclohexyl.

The term "alkenyl" as used herein, and if not specified by the number of carbon atoms, denotes an unsubstituted or substituted hydrocarbon chain radical having from 2 to 8 carbon atoms, preferably from 2 to 4 carbon atoms, and having at least one olefinic double bond, including their different isomers. Examples are vinyl or allyl. A substituted alkenyl group as part of the invention can have the (E) or (Z) configuration. Both isomeric forms of these compounds are embraced by the present invention.

Suitable substituents for the alkenyl group can be selected from 1-2 (preferably 1) substituents selected from cyano, amino-carbonyl and C₁₋₄-alkyl-oxy-carbonyl.

Alkenyl in R² is as defined above, preferably allyl.

Alkenyl for the substituent A is as defined above, preferably a vinyl group or a substituted vinyl group substituted with 1-2 (preferably 1) substituents selected from cyano, amino-carbonyl and C₁₋₄-alkyl-oxy-carbonyl. Examples are vinyl, acrylic acid ethyl ester, acrylonitrile, acrylamide. Both, the (E) or (Z) configuration are embraced by the present invention, however the (E) configuration is preferred.

The term "hydroxy-C₁₋₄-alkyl" as used herein denotes a C₁₋₄-alkyl, preferably a C₁₋₂-alkyl as defined above which is substituted with a hydroxy group. Examples are hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 1-hydroxypropyl, 2-hydroxypropyl, 3-hydroxypropyl, 1-hydroxybutyl, 2-hydroxybutyl, 3-hydroxybutyl, 4-hydroxybutyl.

The term "hydroxy-C₁₋₄-alkyl" for the substituent A is as defined above, preferably hydroxymethyl, 1-hydroxyethyl or 2-hydroxyethyl.

Formula "Z-CH₂-Y-CH₂-" as used herein denotes a chemical group wherein Y represents O or NR wherein R is hydrogen or C₁₋₄-alkyl as defined above, preferably hydrogen; and Z represents heterocyclyl or optionally substituted aryl with 1-3 benzyl-oxy groups (preferably 1-2 benzyl-oxy groups, more preferred 1 benzyl-oxy group). More preferred Z represents aryl or heterocyclyl as defined below, preferably phenyl or 2-pyridyl, 3-pyridyl or 4-pyridyl. Examples of the chemical group of formula "-CH₂-Y-CH₂-Z" are benzylaminomethyl, benzyloxymethyl, (pyridin-4-ylmethyl)aminomethyl, pyridin-4-ylmethoxymethyl, pyridin-3-ylmethoxymethyl, pyridin-4-ylmethoxymethyl or 4-benzyloxy-benzyloxymethyl.

The term "aryl" as used herein denotes an optionally substituted phenyl and naphthyl, both optionally benz-fused to an optionally substituted saturated, partially unsaturated or aromatic monocyclic, bicyclic or tricyclic heterocycle or carbocycle e.g. to cyclohexyl or cyclopentyl.

Suitable substituents for aryl can be selected from 1,2,3,4 or 5 those named for alkyl and C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine or bromine; they can be identical or different from each other. Preferred substituents for aryl are selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine or bromine.

In case more than one substituent is attached to the aryl group, these substituents can be identical or different from each other and are also encompassed by the scope of the present invention. For example dimethoxy-phenyl means that both methoxy substituents may be attached to the phenyl ring in the 2,3-position, the 2,4-position, the 2,5-position, the 2,6-position, the 3,4-position, the 3,5-position and the 3,6-position.

Examples of aryl groups are phenyl, 2-methyl-phenyl, 3-methyl-phenyl, 4-methyl-phenyl, 2,3-dimethylphenyl, 2,4-dimethylphenyl, 2,5-dimethylphenyl, 2,6-dimethylphenyl, 3,4-dimethylphenyl, 3,5-dimethylphenyl, 3,6-dimethylphenyl, 2-methoxy-phenyl, 3-methoxy-phenyl, 4-methoxy-phenyl, 2,3-dimethoxy-phenyl, 2,4-dimethoxy-phenyl, 2,5-dimethoxy-phenyl, 2,6-dimethoxy-phenyl, 3,4-dimethoxy-phenyl, 3,5-dimethoxy-phenyl, 3,6-dimethoxy-phenyl, 2-hydroxyphenyl, 3-hydroxyphenyl, 4-hydroxyphenyl, 2,3-dihydroxyphenyl, 2,4-dihydroxyphenyl, 2,5-dihydroxyphenyl, 2,6-dihydroxyphenyl, 3,4-dihydroxyphenyl, 3,5-dihydroxyphenyl, 3,6-dihydroxyphenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2,3-difluorophenyl, 2,4-difluorophenyl, 2,5-difluorophenyl, 2,6-difluorophenyl, 3,4-difluorophenyl, 3,5-difluorophenyl, 3,6-difluorophenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2,3-dichlorophenyl, 2,4-dichlorophenyl, 2,5-dichlorophenyl, 2,6-dichlorophenyl, 3,4-dichlorophenyl, 3,5-dichlorophenyl, 3,6-dichlorophenyl, 2-bromophenyl, 3-bromophenyl, 4-bromophenyl, 2,3-dibromophenyl, 2,4-dibromophenyl, 2,5-dibromophenyl, 2,6-dibromophenyl, 3,4-dibromophenyl, 3,5-dibromophenyl or 3,6-dibromophenyl.

Aryl in R^1 is as defined above and preferably phenyl, 2-methyl-phenyl, 3-methyl-phenyl, 4-methyl-phenyl, 2,3-dimethylphenyl, 2,4-dimethylphenyl, 2,5-dimethylphenyl, 2,6-dimethylphenyl, 3,4-dimethylphenyl, 3,5-dimethylphenyl, 3,6-dimethylphenyl, 2-chloro-phenyl, 3-chloro-phenyl, 4-chloro-phenyl and most preferably phenyl.

Aryl in the formula " $Z-CH_2-Y-CH_2-$ " as used herein denotes preferably one of the following groups: phenyl, 2-methyl-phenyl, 3-methyl-phenyl, 4-methyl-phenyl, 2,3-dimethylphenyl, 2,4-dimethylphenyl, 2,5-dimethylphenyl, 2,6-dimethylphenyl, 3,4-dimethylphenyl, 3,5-dimethylphenyl, 3,6-dimethylphenyl, 2-chloro-phenyl, 3-chloro-phenyl, 4-chloro-phenyl and most preferably phenyl.

The term "optionally substituted phenyl" in R^3 as used herein denotes a phenyl group which may be substituted with 1,2,3,4 or 5 substituents, preferred 1, 2 or 3 substituents, more preferred 1 or 2 substituents selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine, bromine and cyano or selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine or bromine, preferably methyl, ethyl, methoxy, ethoxy, hydroxy, fluorine, chlorine, bromine; or preferably methyl, ethyl, methoxy, ethoxy,

hydroxy, fluorine, chlorine, bromine and cyano. Further preferred embodiments for "optionally substituted phenyl" in R^3 are 2-substituted phenyl rings and the more preferred 3-substituted phenyl rings. An especially preferred embodiment for "optionally substituted phenyl" in R^3 are 3,5-disubstituted phenyl rings.

5 Preferred examples are phenyl, 2-methyl-phenyl, 3-methyl-phenyl, 4-methyl-phenyl, 2,3-dimethylphenyl, 2,4-dimethylphenyl, 2,5-dimethylphenyl, 2,6-dimethylphenyl, 3,4-dimethylphenyl, 3,5-dimethylphenyl, 3,6-dimethylphenyl, 2-methoxy-phenyl, 3-methoxy-phenyl, 4-methoxy-phenyl, 2,3-dimethoxyphenyl, 2,4-dimethoxyphenyl, 2,5-dimethoxyphenyl, 2,6-dimethoxyphenyl, 3,4-dimethoxyphenyl, 3,5-dimethoxyphenyl, 3,6-dimethoxyphenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2,3-difluorophenyl, 2,4-difluorophenyl, 2,5-difluorophenyl, 2,6-difluorophenyl, 3,4-difluorophenyl, 3,5-difluorophenyl, 3,6-difluorophenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2,3-dichlorophenyl, 2,4-dichlorophenyl, 2,5-dichlorophenyl, 2,6-dichlorophenyl, 3,4-dichlorophenyl, 3,5-dichlorophenyl, 3,6-dichlorophenyl, 2-bromophenyl, 3-bromophenyl, 4-bromophenyl, 2,3-dibromophenyl, 2,4-dibromophenyl, 2,5-dibromophenyl, 2,6-dibromophenyl, 3,4-dibromophenyl, 3,5-dibromophenyl, 3,6-dibromophenyl, 2-cyano-phenyl, 3-cyano-phenyl, 4-cyano-phenyl, 2,3-dicyanophenyl, 2,4-dicyanophenyl, 2,5-dicyanophenyl, 2,6-dicyanophenyl, 3,4-dicyanophenyl, 3,5-dicyanophenyl or 3,6-dicyanophenyl. Most preferred optionally substituted phenyl in R^3 are phenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2,3-dichlorophenyl, 2,4-dichlorophenyl, 2,5-dichlorophenyl, 2,6-dichlorophenyl, 3,4-dichlorophenyl, 3,5-dichlorophenyl, 3,6-dichlorophenyl.

The term "optionally substituted benzyl" in R^3 as used herein denotes a benzyl group which may be substituted with 1,2,3,4 or 5 substituents, preferably 1, 2 or 3 substituents selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine, bromine and cyano or selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine or bromine, preferably methyl, ethyl, methoxy, ethoxy, hydroxy, fluorine, chlorine or bromine. Preferred examples are benzyl, 2-methyl-benzyl, 3-methyl-benzyl, 4-methyl-benzyl, 2,3-dimethylbenzyl, 2,4-dimethylbenzyl, 2,5-dimethylbenzyl, 2,6-dimethylbenzyl, 3,4-dimethylbenzyl, 3,5-dimethylbenzyl, 3,6-dimethylbenzyl, 2-methoxy-benzyl, 3-methoxy-benzyl, 4-methoxy-benzyl, 2,3-dimethoxybenzyl, 2,4-dimethoxybenzyl, 2,5-dimethoxybenzyl, 2,6-dimethoxybenzyl, 3,4-dimethoxybenzyl, 3,5-dimethoxybenzyl, 3,6-dimethoxybenzyl, 2-hydroxy-benzyl, 3-hydroxy-benzyl, 4-hydroxy-benzyl, 2,3-dihydroxybenzyl, 2,4-dihydroxybenzyl, 2,5-dihydroxybenzyl, 2,6-dihydroxybenzyl, 3,4-dihydroxybenzyl, 3,5-dihydroxybenzyl, 3,6-dihydroxybenzyl, 2-fluorobenzyl, 3-fluorobenzyl, 4-fluorobenzyl, 2,3-difluorobenzyl, 2,4-difluorobenzyl, 2,5-difluorobenzyl,

2,6-difluorobenzyl, 3,4-difluorobenzyl, 3,5-difluorobenzyl, 3,6-difluorobenzyl, 2-chlorobenzyl, 3-chlorobenzyl, 4-chlorobenzyl, 2,3-dichlorobenzyl, 2,4-dichlorobenzyl, 2,5-dichlorobenzyl, 2,6-dichlorobenzyl, 3,4-dichlorobenzyl, 3,5-dichlorobenzyl, 3,6-dichlorobenzyl, 2-bromobenzyl, 3-bromobenzyl, 4-bromobenzyl, 2,3-dibromobenzyl, 2,4-dibromobenzyl, 2,5-dibromobenzyl, 2,6-dibromobenzyl, 3,4-dibromobenzyl, 3,5-dibromobenzyl, 3,6-dibromobenzyl, 2-cyano-benzyl, 3-cyano-benzyl, 4-cyano-benzyl, 2,3-dicyanobenzyl, 2,4-dicyanobenzyl, 2,5-dicyanobenzyl, 2,6-dicyanobenzyl, 3,4-dicyanobenzyl, 3,5-dicyanobenzyl or 3,6-dicyanobenzyl. More preferred optionally substituted benzyl in R³ is benzyl, 2-chlorobenzyl, 3-chlorobenzyl, 4-chlorobenzyl, 2,3-dichlorobenzyl, 2,4-dichlorobenzyl, 2,5-dichlorobenzyl, 2,6-dichlorobenzyl, 3,4-dichlorobenzyl, 3,5-dichlorobenzyl, 3,6-dichlorobenzyl and most preferably benzyl is the unsubstituted benzyl group.

In case more than one substituent is attached to the benzyl group, these substituents can be identical or different from each other and are also encompassed by the scope of the present invention. For example dimethoxy-benzyl means that both methoxy substituents may be attached to the phenyl ring in the 2,3-position, the 2,4-position, the 2,5-position, the 2,6-position, the 3,4-position, the 3,5-position and the 3,6-position.

The term "aryl-CH(OH)-" as used herein denotes a hydroxymethyl group which is substituted with an aryl group as defined above. A preferred aryl group is the phenyl group.

The term "heterocyclyl" as used herein denotes optionally substituted aromatic or non-aromatic monocyclic or bicyclic heterocyclic systems which contains one or more hetero atoms selected from nitrogen, oxygen and sulfur, such as furyl, 1-pyrrolyl, 2-pyrrolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 1-indolyl, 2-indolyl or 3-indolyl.

Suitable substituents for heterocyclyl can be selected from 1,2,3,4 or 5 those named for alkyl and C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine or bromine; they can be identical or different from each other. Preferred substituents for heterocyclyl are selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine or bromine.

In case more than one substituent is attached to the heterocyclyl group, these substituents can be identical or different from each other and are also encompassed by the scope of the present invention. For example dimethylpyridyl means that both methyl substituents may be attached to the pyridyl in the chemically possible positions. For example both methyl substituents may be attached to the 2-pyridyl in the 3,4-position, the 4,5-position, the 5,6-position, the 3,5-position, the 3,6-position, the and the 4,6-position.

Both methyl substituents may be attached to the 3-pyridyl in the 2,4-position, the 2,5-position, the 2,6-position, the 4,5-position, the 4,6-position, the and the 5,6-position. Both methyl substituents may be attached to the 4-pyridyl in the 2,3-position, the 2,5-position, the 2,6-position, and the 3,5-position.

5 Examples for "substituted heterocyclyl" are furyl, methylfuryl, dimethylfuryl, ethylfuryl, methoxyfuryl, dimethoxyfuryl, hydroxyfuryl, dihydroxyfuryl, fluorofuryl, difluorofuryl, chlorofuryl, dichlorofuryl, bromofuryl, dibromofuryl, pyrrolyl, methylpyrrolyl, dimethylpyrrolyl, ethylpyrrolyl, methoxypyrrolyl, dimethoxypyrrolyl, hydroxypyrrolyl, dihydroxypyrrolyl, fluoropyrrolyl, difluoropyrrolyl, chloropyrrolyl, 10 dichloropyrrolyl, bromopyrrolyl, dibromopyrrolyl, pyridyl, methylpyridyl, dimethylpyridyl, ethylpyridyl, methoxypyridyl, dimethoxypyridyl, hydroxypyridyl, dihydroxypyridyl, fluoropyridyl, difluoropyridyl, chloropyridyl, dichloropyridyl, bromopyridyl, dibromopyridyl, indolyl, methylindolyl, dimethylindolyl, ethylindolyl, methoxyindolyl, dimethoxyindolyl, hydroxyindolyl, dihydroxyindolyl, fluoroindolyl, 15 difluoroindolyl, chloroindolyl, dichloroindolyl, bromoindolyl or dibromoindolyl. For all the cited examples for "heterocyclyl" these substituents can be at any chemically possible position. For example methylpyridyl means that the methyl substituent may be attached in the 3, 4, 5 or 6 position of a 2-pyridyl or in the 2, 4, 5 or 6 position of a 3-pyridyl or in the 2, 3, 5 or 6 position of a 4-pyridyl.

20 Heterocyclyl in the formula "Z-CH₂-Y-CH₂-" as used herein denoted preferably one of the following groups as defined above, preferably unsubstituted furyl, 1-pyrrolyl, 2-pyrrolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 1-indolyl, 2-indolyl or 3-indolyl and most preferably 2-pyridyl, 3-pyridyl or 4-pyridyl.

 Heterocyclyl in R³ is as defined above and preferably unsubstituted furyl, 1-pyrrolyl, 25 2-pyrrolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 1-indolyl, 2-indolyl or 3-indolyl. More preferably heterocyclyl in R³ is 2-pyridyl, 3-pyridyl or 4-pyridyl and most preferred 4-pyridyl.

 The term "C₁₋₄-alkyl substituted with optionally substituted heterocyclyl" as used herein denotes a C₁₋₄-alkyl, preferably C₁₋₂-alkyl group as defined above which may be 30 substituted with an optionally substituted heterocyclyl group, preferably with an optionally substituted pyridyl group as defined above which may be substituted with 1,2,3 or 4 substituents, preferably 1 or 2 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine or bromine, preferably methyl, ethyl, methoxy, ethoxy, hydroxy, fluorine, chlorine or bromine. Examples are furylmethyl, furylethyl, furylpropyl, 35 furylbutyl, methylfurylmethyl, methylfurylethyl, dimethylfurylmethyl, ethylfurylmethyl,

methoxyfurylmethyl, methoxyfurylethyl, dimethoxyfurylmethyl, hydroxyfurylmethyl, hydroxyfurylethyl, dihydroxyfurylmethyl, fluorofurylmethyl, difluorofurylmethyl, chlorofurylmethyl, chlorofurylethyl, dichlorofurylmethyl, dichlorofurylethyl, bromofurylmethyl, dibromofurylmethyl, pyrrolylmethyl, pyrrolylethyl, pyrrolylpropyl, pyrrolylbutyl, methylpyrrolylmethyl, methylpyrrolylethyl, dimethylpyrrolylmethyl, ethylpyrrolylmethyl, methoxypyrrolylmethyl, methoxypyrrolylethyl, dimethoxypyrrolylmethyl, hydroxypyrrolylmethyl, hydroxypyrrolylethyl, dihydroxypyrrolylmethyl, fluoropyrrolylmethyl, difluoropyrrolylmethyl, chloropyrrolylmethyl, chloropyrrolylethyl, dichloropyrrolylmethyl, dichloropyrrolylmethyl, bromopyrrolylmethyl, dibromopyrrolylmethyl, pyridylmethyl (more specific examples are 2-pyridylmethyl, 3-pyridylmethyl, 4-pyridylmethyl), pyridylethyl, pyridylpropyl, pyridylbutyl, methylpyridylmethyl, methylpyridylethyl, dimethylpyridylmethyl, ethylpyridylmethyl, methoxypyridylmethyl, methoxypyridylethyl, dimethoxypyridylmethyl, hydroxypyridylmethyl, hydroxypyridylethyl, dihydroxypyridylmethyl, fluoropyridylmethyl, difluoropyridylmethyl, chloropyridylmethyl, chloropyridylethyl, dichloropyridylmethyl, dichloropyridylmethyl, bromopyridylmethyl, dibromopyridylmethyl, indolylmethyl, indolylethyl, indolylpropyl, indolylbutyl, methylindolylmethyl, methylindolylethyl, dimethylindolylmethyl, ethylindolylmethyl, methoxyindolylmethyl, methoxyindolylethyl, dimethoxyindolylmethyl, hydroxyindolylmethyl, hydroxyindolylethyl, dihydroxyindolylmethyl, fluoroindolylmethyl, difluoroindolylmethyl, chloroindolylmethyl, chloroindolylethyl, dichloroindolylmethyl, dichloroindolylmethyl, bromoindolylmethyl or dibromoindolylmethyl.

For all the cited examples for "heterocyclyl" these substituents can be at any chemically possible position. For example methylpyridyl means that the methyl substituent may be attached in the 3, 4, 5 or 6 position of a 2-pyridyl or in the 2, 4, 5 or 6 position of a 3-pyridyl or in the 2, 3, 5 or 6 position of a 4-pyridyl.

The term "X" represents S and O, preferably S.

The term halogen stands for fluorine, chlorine, bromine and iodine.

30

Any functional (i.e. reactive) group present in a side-chain may be protected, with the protecting group being a group which is known per se, for example, as described in "Protective Groups in Organic Synthesis", 2nd Ed., T.W. Greene and P.G.M. Wuts, John

Wiley & Sons, New York, NY, 1991. For example, an amino group can be protected by tert.-butoxycarbonyl (BOC) or benzyloxycarbonyl (Z).

The compounds of this invention may contain one or more asymmetric carbon atoms and may therefore occur as racemates and racemic mixtures, single enantiomers, 5 diastereomeric mixtures and individual diastereomers. Furthermore, where a compound of the invention contains an olefinic double bond, this can have the (E) or (Z) configuration. Also, each chiral center may be of the R or S configuration. All such isomeric forms of these compounds are embraced by the present invention.

Compounds of formula I which are acidic can form pharmaceutically acceptable 10 salts with bases such as alkali metal hydroxides, e.g. sodium hydroxide and potassium hydroxide; alkaline earth metal hydroxides, e.g. calcium hydroxide, barium hydroxide and magnesium hydroxide, and the like; with organic bases e.g. N-ethyl piperidine, dibenzylamine, and the like. Those compounds of formula (I) which are basic can form pharmaceutically acceptable salts with inorganic acids, e.g. with hydrohalic acids such as 15 hydrochloric acid and hydrobromic acid, sulphuric acid, nitric acid and phosphoric acid, and the like, and with organic acids, e.g. with acetic acid, tartaric acid, succinic acid, fumaric acid, maleic acid, malic acid, salicylic acid, citric acid, methanesulphonic acid and p-toluene sulphonic acid, and the like. The formation and isolation of such salts can be carried out according to methods known in the art.

20

Preferred embodiments of the invention are compounds of formula I wherein

R^1 is hydrogen, C_{1-12} -alkyl, C_{3-8} -cycloalkyl, aryl or C_{1-4} -alkyl substituted with phenyl optionally substituted with 1-5 substituents selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine and bromine,

25

preferred wherein

R^1 is C_{1-12} -alkyl, aryl or C_{1-4} -alkyl substituted with phenyl optionally substituted with 1-5 substituents selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine and bromine,

more preferred wherein

30

R^1 is C_{1-7} -alkyl, phenyl or C_{1-4} -alkyl substituted with phenyl,

most preferred wherein

R^1 is C_{1-7} -alkyl or C_{1-4} -alkyl substituted with phenyl;

R^2 is hydrogen, C_{2-4} -alkenyl, C_{1-12} -alkyl or C_{1-4} -alkyl substituted with phenyl optionally substituted with 1-5 substituents selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine and bromine,

5

preferred wherein

R^2 is C_{2-4} -alkenyl, C_{1-12} -alkyl or C_{1-4} -alkyl substituted with phenyl optionally substituted with 1-5 substituents selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine and bromine,

10

more preferred wherein

R^2 is C_{2-4} -alkenyl, C_{1-7} -alkyl or C_{1-4} -alkyl substituted with phenyl optionally substituted with 1-5 substituents selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine and bromine,

most preferred wherein

15

R^2 is C_{2-4} -alkenyl, C_{1-7} -alkyl or C_{1-4} -alkyl substituted with phenyl;

R^3 is C_{1-12} -alkyl, C_{3-8} -cycloalkyl, phenyl, benzyl or heterocyclyl,

wherein a phenyl ring may be substituted with 1-5 substituents selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine, bromine and cyano,

20

preferred wherein

R^3 is C_{1-12} -alkyl, C_{3-8} -cycloalkyl, phenyl, benzyl or heterocyclyl,

wherein a phenyl ring may be substituted with 1-5 substituents selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine, bromine and cyano,

25

more preferred wherein

R^3 is C_{1-7} -alkyl, C_{3-6} -cycloalkyl, phenyl, benzyl or heterocyclyl,

wherein a phenyl ring may be substituted with 1-5 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine, bromine and cyano,

most preferred wherein

5 R³ is C₁₋₄-alkyl, C₃₋₆-cycloalkyl or optionally substituted phenyl,

wherein phenyl may be substituted with 1-5 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine, bromine and cyano;

10 A is C₁₋₁₂-alkyl, hydroxy-C₁₋₄-alkyl, C₂₋₄-alkenyl, substituted C₂₋₄-alkenyl, aryl-CH(OH)-, C₁₋₄-alkyl substituted with C₁₋₄-alkoxy, C₁₋₄-alkyl-carbonyl-oxy-, heterocyclyl-oxy or A is C₁₋₄-alkyl substituted with phenyl optionally substituted with 1-5 substituents or with heterocyclyl optionally substituted with 1-4 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine,

15 wherein substituted C₂₋₄-alkenyl is substituted with 1-2 substituents selected from cyano, amino-carbonyl and C₁₋₄-alkyl-oxy-carbonyl, or

A signifies a group of formula Z-CH₂-Y-CH₂-,

wherein Y represents O or NR wherein R is hydrogen or C₁₋₄-alkyl and

Z represents heterocyclyl or optionally substituted aryl with 1-3 benzyl-oxy groups,

20 preferred wherein

A is C₁₋₇-alkyl, hydroxy-C₁₋₄-alkyl, C₂₋₄-alkenyl, substituted C₂₋₄-alkenyl, aryl-CH(OH)-, C₁₋₄-alkyl substituted with C₁₋₄-alkoxy, C₁₋₄-alkyl-carbonyl-oxy-, heterocyclyl-oxy or A is C₁₋₄-alkyl substituted with phenyl optionally substituted with 1-5 substituents or with heterocyclyl optionally substituted with 1-4 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine,

25

wherein substituted C₂₋₄-alkenyl is substituted with 1-2 substituents selected from cyano, amino-carbonyl and C₁₋₄-alkyl-oxy-carbonyl, or

A signifies a group of formula $Z-CH_2-Y-CH_2-$,

wherein Y represents O or NR wherein R is hydrogen or C_{1-4} -alkyl and

Z represents heterocyclyl or optionally substituted aryl with 1-3 benzyl-oxy groups,

5 more preferred wherein

A is C_{1-7} -alkyl, hydroxy- C_{1-4} -alkyl, C_{2-4} -alkenyl, substituted C_{2-4} -alkenyl, phenyl-CH(OH)-, C_{1-4} -alkyl substituted with C_{1-4} -alkoxy, C_{1-4} -alkyl-carbonyl-oxy-, heterocyclyl-oxy or A is C_{1-4} -alkyl substituted with phenyl optionally substituted with 1-5 substituents or with pyridyl optionally substituted with 1-4 substituents
10 selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine and bromine,

wherein substituted C_{2-4} -alkenyl is substituted with 1-2 substituents selected from cyano, amino-carbonyl and C_{1-4} -alkyl-oxy-carbonyl, or

A signifies a group of formula $Z-CH_2-Y-CH_2-$,

wherein Y represents O or NR wherein R is hydrogen or C_{1-4} -alkyl and

15 Z represents pyridyl or optionally substituted phenyl with 1-3 benzyl-oxy groups,

most preferred wherein

A is C_{1-7} -alkyl, C_{2-4} -alkenyl, substituted C_{2-4} -alkenyl, C_{1-4} -alkyl substituted with phenyl optionally substituted with 1-5 substituents or with pyridyl optionally substituted with 1-4 substituents selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine and bromine,
20

wherein substituted C_{2-4} -alkenyl is substituted with 1-2 substituents selected from cyano, amino-carbonyl and C_{1-4} -alkyl-oxy-carbonyl, or

A signifies a group of formula $Z-CH_2-Y-CH_2-$,

25 wherein Y represents O or NR wherein R is hydrogen or C_{1-4} -alkyl and

Z represents pyridyl or optionally substituted phenyl with 1-3 benzyl-oxy groups;

X represents S or O;

with the provisos that

- 5 (i) only one of R^1 and R^2 is hydrogen;
(ii) when X is O, then R^1 cannot be aryl;

as well as hydrolyzable esters or ethers of compounds of formula I and pharmaceutically acceptable salts thereof.

- 10 Further preferred embodiments of the invention are compounds of formula I wherein

R^1 is C_{1-7} -alkyl,

preferred wherein

R^1 is C_{1-4} -alkyl,

- 15 more preferred wherein

R^1 is isopropyl;

R^2 is C_{1-7} -alkyl,

preferred wherein

R^2 is C_{1-4} -alkyl,

- 20 more preferred wherein

R^2 is methyl or ethyl,

most preferred wherein

R^2 is methyl;

R^3 is optionally substituted phenyl,

wherein phenyl may be substituted with 1-5 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine, bromine and cyano,

preferred wherein

R³ is optionally substituted phenyl,

5 wherein phenyl may be substituted with 1-3 substituents selected from C₁₋₂-alkyl, C₁₋₂-alkoxy, hydroxy, fluorine, chlorine, bromine and cyano,

more preferred wherein

R³ is optionally substituted phenyl,

 wherein phenyl may be substituted with 1-3 chlorine substituents;

10 most preferred wherein

R³ is optionally substituted phenyl,

 wherein phenyl may be substituted with 1-2 chlorine substituents;

A is C₁₋₄-alkyl substituted with phenyl optionally substituted with 1-3 substituents or
with pyridyl optionally substituted with 1-2 substituents selected from C₁₋₄-alkyl,
15 C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine, or

A signifies a group of formula Z-CH₂-Y-CH₂-,

 wherein Y represents O or NR wherein R is hydrogen or C₁₋₄-alkyl and

 Z represents pyridyl or phenyl,

preferred wherein

20 A is C₁₋₂-alkyl substituted with phenyl optionally substituted with 1-3 substituents or
with pyridyl optionally substituted with 1-2 substituents selected from C₁₋₄-alkyl,
C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine, or

A signifies a group of formula Z-CH₂-Y-CH₂-,

 wherein Y represents O or NR wherein R is hydrogen or C₁₋₂-alkyl and

Z represents pyridyl or phenyl,

more preferred wherein

A is C₁₋₂-alkyl substituted with phenyl or pyridyl, or

A signifies a group of formula Z-CH₂-Y-CH₂-,

5 wherein Y represents O or NR wherein R is hydrogen or C₁₋₂-alkyl and

Z represents pyridyl or phenyl,

most preferred wherein

A is C₁₋₂-alkyl substituted with pyridyl, or

A signifies a group of formula Z-CH₂-Y-CH₂-,

10 wherein Y represents O or NR wherein R is hydrogen or C₁₋₂-alkyl and

Z represents pyridyl;

X represents S or O;

as well as hydrolyzable esters or ethers of compounds of formula I and pharmaceutically acceptable salts thereof.

15

Another embodiment of the invention are compounds of formula I wherein

R¹ is hydrogen, C₁₋₁₂-alkyl, C₃₋₈-cycloalkyl, aryl or C₁₋₄-alkyl substituted with phenyl optionally substituted with 1-5 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine;

20 R² is hydrogen, C₁₋₁₂-alkyl or C₁₋₄-alkyl substituted with phenyl optionally substituted with 1-5 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine;

R³ is C₁₋₁₂-alkyl, C₃₋₈-cycloalkyl, phenyl or benzyl,

wherein a phenyl ring may be substituted with 1-5 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine;

A is hydroxy-C₁₋₄-alkyl or aryl-CH(OH)-; or

A signifies a group of formula Z-CH₂-Y-CH₂-,

5 wherein Y represents O or NR wherein R is hydrogen or C₁₋₄-alkyl and

 Z represents aryl or heterocyclyl; or

A signifies C₁₋₄-alkyl substituted with phenyl optionally substituted with 1-5 substituents or with heterocyclyl optionally substituted with 1-4 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine;

10 X represents S or O;

with the proviso that only one of R¹ and R² is hydrogen;

as well as hydrolyzable esters or ethers of compounds of formula I and pharmaceutically acceptable salts thereof.

15 An especially preferred embodiment of the invention are compounds of formula I wherein






 X represents S;







as well as hydrolyzable esters or ethers of compounds of formula I and pharmaceutically acceptable salts thereof.






More preferred embodiments of compounds of formula I, as well as hydrolyzable esters or ethers of compounds of formula I and pharmaceutically acceptable salts thereof, are listed in table 1:







5







Table 1

STRUCTURE	SYSTEMATIC NAME
	4-(3,5-Dichlorophenylthio)-5-hydroxymethyl-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone
	5-(Benzylaminomethyl)-4-(3,5-dichlorophenylthio)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone
	4-(3,5-Dichlorophenylthio)-5-(alpha(RS)-hydroxybenzyl)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone
	4-(3,5-Dichlorophenylthio)-5-hydroxymethyl-1-methyl-3-phenyl-1,3-dihydro-2-imidazolone
	5-Benzyloxymethyl-4-(3,5-dichlorophenylthio)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone








	<p>4-(3,5-Dichlorophenylthio)-3-isopropyl-1-methyl-5-[[pyridin-4-ylmethyl]-amino]-methyl-1,3-dihydro-2-imidazolone</p>
	<p>5-(Benzylaminomethyl)-3-isopropyl-1-methyl-4-phenylthio-1,3-dihydro-2-imidazolone</p>
	<p>3-Isopropyl-1-methyl-4-phenylthio-5-[[pyridin-4-ylmethyl]amino]methyl-1,3-dihydro-2-imidazolone</p>
	<p>5-Benzyloxymethyl-4-(3,5-dichlorophenoxy)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone</p>
	<p>5-Benzyloxymethyl-4-(3,5-dichlorophenylthio)-1-methyl-3-phenyl-1,3-dihydro-2-imidazolone</p>
	<p>4-(3,5-Dichlorothio)-3-isopropyl-1-methyl-5-(pyridin-4-ylmethoxymethyl)-1,3-dihydro-2-imidazolone</p>









	5-(Benzylaminomethyl)-4-(3,5-dimethylphenylthio)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone
	4-Cyclohexylthio-3-isopropyl-1-methyl-5-[[pyridin-4-ylmethyl]amino]methyl-1,3-dihydro-2-imidazolone
	4-(3,5-Dimethylphenylthio)-3-isopropyl-1-methyl-5-[[pyridin-4-ylmethyl]amino]methyl-1,3-dihydro-2-imidazolone
	4-Benzylthio-3-isopropyl-1-methyl-5-[[pyridin-4-ylmethyl]amino]methyl-1,3-dihydro-2-imidazolone
	4-(3-Chlorophenylthio)-3-isopropyl-1-methyl-5-[[pyridin-4-ylmethyl]amino]methyl-1,3-dihydro-2-imidazolone





	3-Isopropyl-4-isopropylthio-1-methyl-5-[[pyridin-4-ylmethyl]amino]methyl-1,3-dihydro-2-imidazolone
	4-Isobutylthio-3-isopropyl-1-methyl-5-[[pyridin-4-ylmethyl]amino]methyl-1,3-dihydro-2-imidazolone
	4-(3,5-Dichlorophenylthio)-3-isopropyl-1-(4-methoxybenzyl)-5-[[pyridin-4-ylmethyl]amino]methyl-1,3-dihydro-2-imidazolone
	4-(3,5-Dichlorophenylthio)-5-hydroxymethyl-3-isopropyl-1-(4-methoxybenzyl)-1,3-dihydro-2-imidazolone
	4-(3,5-Dichlorophenylthio)-3-isopropyl-1-methyl-5-phenethyl-1,3-dihydro-2-imidazolone
	4-(3,5-Dichlorophenylthio)-5-(2-hydroxybenzyl)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone

	<p>3-Benzyl-4-(3,5-dichlorophenylthio)-1-methyl-5- {[(pyridin-4-ylmethyl)amino]methyl}-1,3-dihydro-2- imidazolone</p>
	<p>1,3-Dibenzyl-4-(3,5-dichlorophenylthio)-5- {[(pyridin-4-ylmethyl)amino]methyl}-1,3-dihydro-2-imidazolone</p>
	<p>1-Benzyl-4-(3,5-dichlorophenylthio)-3-isopropyl-5- {[(pyridin-4-ylmethyl)amino]methyl}-1,3-dihydro-2- imidazolone</p>
	<p>1-Benzyl-4-(3,5-dichlorophenylthio)-3-propyl-5- {[(pyridin-4-ylmethyl)amino]methyl}-1,3-dihydro-2- imidazolone</p>
	<p>5-Benzyl-4-(3,5-dichlorophenylthio)-3-isopropyl-1- methyl-1,3-dihydro-2-imidazolone</p>
	<p>3-Benzyl-4-(3,5-dichlorophenylthio)-5-hydroxymethyl- 1-methyl-1,3-dihydro-2-imidazolone</p>

	<p>4-(3,5-Dichlorophenoxy)-3-isopropyl-1-methyl-5- {[(pyridin-4-ylmethyl)amino]methyl}-1,3-dihydro-2- imidazolone</p>
	<p>4-sec-Butylthio-3-isopropyl-1-methyl-5- {[(pyridin-4-ylmethyl)amino]methyl}-1,3-dihydro-2- imidazolone</p>
	<p>3-Isopropyl-4-(3-methoxyphenylthio)-1-methyl-5- {[(pyridin-4-ylmethyl)amino]methyl}-1,3-dihydro-2- imidazolone</p>
	<p>3-Benzyl-4-(3,5-dichlorophenylthio)-5-(1-hydroxyethyl)- 1-methyl-1,3-dihydro-2-imidazolone</p>
	<p>4-(3,5-Dichloro-phenylsulfanyl)-3-isopropyl-5- methoxymethyl-1-methyl-1,3-dihydro-imidazol-2-one</p>
	<p>4-Hydroxymethyl-3-methyl-1-phenyl-5-(pyridin-4- ylsulfanyl)-1,3-dihydro-imidazol-2-one</p>

	<p>1-Benzyl-4-hydroxymethyl-3-methyl-5-(pyridin-4-ylsulfanyl)-1,3-dihydro-imidazol-2-one; hydrochloric acid</p>
	<p>4-(3,5-Dichloro-phenylsulfanyl)-3-isopropyl-1-methyl-5-(pyridin-3-ylmethoxymethyl)-1,3-dihydro-imidazol-2-one</p>
	<p>4-(4-Benzyloxy-benzyloxymethyl)-5-(3,5-dichloro-phenylsulfanyl)-1-isopropyl-3-methyl-1,3-dihydro-imidazol-2-one</p>
	<p>Acetic acid 5-(3,5-dichloro-phenylsulfanyl)-1-isopropyl-3-methyl-2-oxo-2,3-dihydro-1H-imidazol-4-ylmethyl ester</p>
	<p>4-(3,5-Dichloro-phenylsulfanyl)-3-isopropyl-1-methyl-5-(pyridin-4-yloxymethyl)-1,3-dihydro-imidazol-2-one</p>
	<p>3-(5-Benzyl-3-isopropyl-1-methyl-2-oxo-2,3-dihydro-1H-imidazol-4-ylsulfanyl)-benzonitrile</p>
	<p>4-Benzyl-5-isobutylsulfanyl-1-isopropyl-3-methyl-1,3-dihydro-imidazol-2-one</p>

	4-Benzyl-1-isopropyl-5-isopropylsulfanyl-3-methyl-1,3-dihydro-imidazol-2-one
	4-Benzyl-1-isopropyl-3-methyl-5-methylsulfanyl-1,3-dihydro-imidazol-2-one
	4-(3,5-Dichloro-phenylsulfanyl)-5-isobutyl-3-isopropyl-1-methyl-1,3-dihydro-imidazol-2-one
	4-Benzyl-5-(3,5-dichloro-phenylsulfanyl)-3-ethyl-1-isopropyl-1,3-dihydro-imidazol-2-one
	1-Allyl-5-benzyl-4-(3,5-dichloro-phenylsulfanyl)-3-isopropyl-1,3-dihydro-imidazol-2-one
	4-(3,5-Dichloro-phenylsulfanyl)-1-ethyl-3-isopropyl-5-pyridin-4-ylmethyl-1,3-dihydro-imidazol-2-one
	1-Allyl-4-(3,5-dichloro-phenylsulfanyl)-3-isopropyl-5-pyridin-4-ylmethyl-1,3-dihydro-imidazol-2-one
	[E]-3-[5-(3,5-Dichloro-phenylsulfanyl)-1-isopropyl-3-methyl-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-acrylic acid ethyl ester

	3-[5-(3,5-Dichloro-phenylsulfanyl)-1-isopropyl-3-methyl-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-acrylonitrile
	3-[5-(3,5-Dichloro-phenylsulfanyl)-1-isopropyl-3-methyl-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-acrylamide
	1-Benzyl-5-(3,5-dichloro-phenylsulfanyl)-3-methyl-4-vinyl-1,3-dihydro-imidazol-2-one
	4-(3,5-Dichloro-phenylsulfanyl)-3-isopropyl-1-methyl-5-pyridin-4-ylmethyl-1,3-dihydro-imidazol-2-one

The imidazolone derivatives provided by the present invention are useful in the treatment of the human or animal body.

The imidazolone derivatives provided by the present invention are inhibitors of the human immunodeficiency virus reverse transcriptase enzyme. Accordingly, the present
5 imidazolone derivatives are therapeutically active substances in the treatment of diseases mediated by the human immunodeficiency virus (HIV) and can be used as medicaments for the treatment of such diseases.

They can be used as medicaments, especially for treating viral diseases, immune
10 mediated conditions or diseases, bacterial diseases, parasitic diseases, inflammatory diseases, hyperproliferative vascular diseases, tumors, and cancer.


In particular, compounds of the present invention and pharmaceutical compositions containing the same are useful as chemotherapeutic agents, inhibitors of viral replication and modulators of the immune system, and can be used for the treatment of diseases
15 mediated by the human immunodeficiency virus (HIV) other viral diseases such as retroviral infections (either alone or in combination with other antiviral agents such as interferon or derivatives thereof, such as conjugates with polyethylene glycol).

They can be used alone, or in combination with other therapeutically active agents, for example, an immunosuppressant, a chemotherapeutic agent, an anti-viral agent, an
20 antibiotic, an anti-parasitic agent, an anti-inflammatory agent, an anti-fungal agent and/or an anti-vascular hyperproliferation agent.

Compounds, whenever prepared by the processes of the present invention are also an object of the present invention.


The compounds of the present invention can be prepared as shown in the following scheme:

Reaction scheme 1:



5 wherein R^1 , R^2 , R^3 , X and Z are as defined for compounds of formula I.


In accordance with the present invention, the preparation of compounds of formula



wherein R^1 , R^2 , R^3 and X are as defined in formula I;

10 comprises

reacting chloro compound of formula VII



wherein R¹ and R² are as defined in formula I

with nucleophilic agent R³XH

5 wherein R³ and X are as defined in formula I.

The reaction is described in more detail in reaction scheme 1 (fifth reaction step).

In reaction scheme 1, the first reaction step is carried out in that glycine derivative of
 10 formula II (commercially available or prepared according to known methods in the art for
 example alkylation reaction of the nitrogen function of glycine) is reacted with an
 isocyanate derivative of formula III (commercially available or synthesised according to
 known methods from textbooks on organic chemistry e.g. from J. March (1992),
 "Advanced Organic Chemistry: Reactions, Mechanisms and Structure", 4th ed. John Wiley
 15 and Sons) in an appropriate solvent, in the presence of 0.5 – 1.5 equivalents, preferably 0.7
 – 1.3 equivalents, most preferred 0.9 equivalents of a base such as potassium carbonate,
 sodium carbonate, magnesium carbonate, calcium carbonate, potassium hydroxide,
 sodium hydroxide, magnesium hydroxide, calcium hydroxide, N(CH₃)₃, N(C₂H₅)₃,
 N(n-C₃H₇)₃, N(iso-C₃H₇)₃, preferably sodium hydroxide. The reaction is conveniently
 20 carried out at a reaction temperature from -5°C to boiling temperature of the reaction
 mixture, preferably at a reaction temperature between 0°C and about 80°C. The reaction
 can optionally be carried out under an inert atmosphere such as nitrogen or argon
 atmosphere. An appropriate solvent may be water or an organic solvent such as alcohols
 (e.g. methanol, ethanol, propanol, butanol, octanol or cyclohexanol), ethers (e.g.
 25 tetrahydrofuran, diethyl ether, dibutyl ether or dioxane), halogenated hydrocarbons (e.g.
 dichloromethane or trichloromethane), hydrocarbons (e.g. cyclohexane, methyl
 cyclohexane, decaline, benzene, toluene, o-xylene, m-xylene or p-xylene) or a mixtures of
 the mentioned solvents, preferably water. After the reaction, the system is neutralised with

an acid, preferably a protic acid such as HBr, HI, AcOH or citric acid, most preferred HCl, to yield the urea of formula IV.

In the second step of the reaction, urea of formula IV is converted in a ring-closure reaction to imidazoldione derivatives of formula V by dissolving compound of formula IV
5 in water and an acid, preferably a protic acid such as HCl, HBr, HI, AcOH, citric or other, most preferred HCl. The reaction is conveniently carried out at a reaction temperature from room temperature to boiling temperature of the reaction mixture, preferably at a reaction temperature from 60°C to boiling temperature of the reaction mixture. The reaction can optionally be carried out in an inert organic solvent such as ethers (e.g.
10 tetrahydrofuran, diethyl ether, dibutyl ether or dioxane), polar aprotic solvents such as dimethylsulfoxide (DMSO) or dimethylacetamide, halogenated hydrocarbons such as dichloromethane or trichloromethane, hydrocarbons such as cyclohexane, methyl cyclohexane, decaline, benzene, toluene, o-xylene, m-xylene or p-xylene, or mixtures of the mentioned solvents.


In the third step of the reaction, imidazoldione derivatives of formula V is reacted with di-C₁₋₄-alkylacetal of dimethylformamide, preferably dimethylformamide
15 dimethylacetal (commercially available or synthesized according to methods known from textbooks about organic chemistry e.g. from J. March (1992), "Advanced Organic Chemistry: Reactions, Mechanisms, and Structure", 4th ed. John Wiley & Sons) to attach a
20 dimethylaminomethylene group to the imidazoldione ring and to obtain the corresponding compound of formula VI. The reaction is carried out at a reaction temperature from room temperature to boiling temperature of the reaction mixture, preferably at a reaction temperature from 60°C to boiling temperature of the reaction mixture, most preferred at a reaction temperature from 80°C to boiling temperature of the
25 reaction mixture. The reaction can optionally be carried out in an inert organic solvent such as ethers (e.g. tetrahydrofuran, diethyl ether, dibutyl ether or dioxane), polar aprotic solvents such as dimethylsulfoxide (DMSO), halogenated hydrocarbons such as dichloromethane or trichloromethane, hydrocarbons such as cyclohexane, methyl cyclohexane, decaline, benzene, toluene, o-xylene, m-xylene or p-xylene, or mixtures of
30 the mentioned solvents.

In the fourth step of the reaction, dimethylaminomethylene imidazoldione derivatives of formula VI are converted to the corresponding carbaldehyde chloro
imidazolone compound of formula VII as described in US 4,345,936 or known from
textbooks about organic chemistry. The reaction which includes a
35 dimethylaminomethylene/carbaldehyde and oxo/chlorine exchange may be carried out in

the presence of a disubstituted formamide such as N,N-dimethylformamide, N,N-methylphenylformamide or N,N-diphenylformamide, preferably dimethylformamide in the presence of POCl_3 according the Vilsmeier reaction or in the presence of other chlorinating agents such as PCl_5 , SOCl_2 or $(\text{COCl})_2$. The reaction is optionally carried out
5 under an inert atmosphere such as nitrogen or argon atmosphere at a reaction temperature from room temperature to boiling temperature of the reaction mixture, preferably at a reaction temperature between about 50°C and about 150°C . The reaction can optionally be carried out in an inert organic solvent such as ethers (e.g. tetrahydrofuran, diethyl ether, dibutyl ether or dioxane), polar aprotic solvents such as
10 dimethylsulfoxide (DMSO) or N,N-dimethylacetamide, halogenated hydrocarbons such as dichloromethane or trichloromethane, hydrocarbons such as cyclohexane, methyl cyclohexane, decaline, benzene, toluene, o-xylene, m-xylene or p-xylene, or a mixtures of the mentioned solvents.

In the fifth step of the reaction, chloro compounds of formula VII are reacted with
15 nucleophilic thiol R^3SH or nucleophilic alcohol R^3OH (both agents are commercially available or can be synthesized according to methods known from textbooks about organic chemistry e.g. from J. March (1992), "Advanced Organic Chemistry: Reactions, Mechanisms, and Structure", 4th ed. John Wiley & Sons, to obtain the corresponding compound of formula VIII. The reaction is carried out in an appropriate solvent in the
20 presence of a suitable base such as n-BuLi, sodium hydride, trialkylamine (e.g. trimethylamine or triethylamine), potassium carbonate, sodium carbonate, magnesium carbonate, calcium carbonate, preferably triethylamine, N-methyl morpholine or potassium carbonate. The reaction is conveniently carried out under an inert atmosphere such as nitrogen or argon atmosphere at a reaction temperature from 0°C to boiling
25 temperature of the reaction mixture, preferably at a reaction temperature between about 10°C and about 60°C . Appropriate solvents for the reaction are ethers (e.g. tetrahydrofuran, diethyl ether, dibutyl ether or dioxane), polar aprotic solvents such as dimethylsulfoxide (DMSO) or N,N-dimethylacetamide, halogenated hydrocarbons such as dichloromethane or trichloromethane, hydrocarbons such as cyclohexane, methyl
30 cyclohexane, decaline, benzene, toluene, o-xylene, m-xylene or p-xylene, or a mixtures of the mentioned solvents, preferably dichloromethane or trichloromethane.

Also part of the present invention are novel compounds of formula VIII-a



wherein R^1 , R^2 and X are as defined in formula I and R^3 is C_{7-12} -alkyl, C_{3-8} -cycloalkyl, optionally substituted phenyl or optionally substituted benzyl

5 wherein phenyl or benzyl may be substituted with 1-5 substituents selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine or bromine

In the sixth step of the reaction, the aldehyde of formula VIII is reduced in the presence of a reducing agent to obtain the corresponding alcohol derivative of formula Ia. Reducing agents conveniently used for the reaction are preferably sodium borohydride or
 10 other reducing agents such as lithium borohydride, sodium triacetoxyborohydride, hydrogen over a catalyst or reducing agents known in the art applied according to known methods described in textbooks on organic chemistry e.g. from J. March (1992), "Advanced Organic Chemistry: Reactions, Mechanisms and Structure", 4th ed. John Wiley and Sons. The reduction reaction is conveniently carried out in an organic solvent for
 15 example alcoholic solvents such as methanol, ethanol, propanol, butanol, octanol or cyclohexanol, preferably methanol or ethanol or ethers such as tetrahydrofuran, diethyl ether, dibutyl ether, dioxane or diglyme, preferably tetrahydrofuran or a mixture of the mentioned solvents such as methanol and tetrahydrofuran or ethanol and tetrahydrofuran. The reaction is carried out at a reaction temperature between about -10°C and about 60°C ,
 20 preferably at room temperature. The reduction reaction can also be carried out as described in textbooks about organic chemistry e.g. from J. March (1992), "Advanced Organic Chemistry: Reactions, Mechanisms, and Structure", 4th ed. John Wiley & Sons.

In the seventh step of the reaction, the alcohol derivative of formula Ia is derivatized to the corresponding ether derivative of formula Ib. The derivatisation reaction is
 25 conveniently carried out with an electrophilic agent $\text{L-CH}_2\text{-Z}$, wherein Z is as defined for compounds of formula I and L is a leaving group such as bromide, chloride, iodide, tosylate, triflate or any other according to methods known from textbooks on organic chemistry e.g. from J. March (1992), "Advanced Organic Chemistry: Reactions, Mechanisms and Structure", 4th ed. John Wiley and Sons). The reaction is carried out as

described in the literature, for example in the presence of a base such as sodium hydride, lithium hydride, potassium carbonate or triethylamine in an appropriate organic solvent such as tetrahydrofuran (THF) or polar aprotic solvents like dimethylsulfoxide (DMSO), N,N-dimethylacetamide or N,N-dimethylformamide (DMF), preferably DMF or THF, at a
5 reaction temperature between about -10°C and about 60°C, preferably at room temperature.

For the synthesis of compounds of formula Ia or Ib wherein R^1 , R^3 , X and Z are as defined for compounds of formula I and R^2 is hydrogen, the corresponding compound of formula VI is N-protected with an appropriate protecting group for example
10 4-methoxybenzyl, benzyl or 3,4-dimethoxybenzyl and then further reacted according to the reaction steps 4-6 (and optionally 7), and finally deprotected according to methods known from textbooks on protecting chemistry ('Protecting Groups in Organic Synthesis', second edition, Theodora W. Greene and Peter G. M. Wuts, Wiley-Interscience, 1991) for example in the presence of hydrogen and appropriate catalyst such as Pd/C or Pt/C.
15 Compounds of formula Ia or Ib wherein R^1 , R^3 , X and Z are as defined for compounds of formula I and R^2 is hydrogen are obtained. The protecting reaction is for example carried out in a polar aprotic solvents like dimethylsulfoxide (DMSO), dimethylacetamide or N,N-dimethylformamide (DMF), preferably dimethylformamide in the presence of a base such as lithium hydride or sodium hydride, preferably sodium hydride at a reaction
20 temperature between about -10°C and about 60°C, preferably at room temperature. The reaction system is then treated with an appropriate protecting agent such as 4-methoxybenzyl chloride, 4-methoxybenzyl bromide, benzyl chloride, benzyl bromide, 3,4-dimethoxybenzyl chloride or 3,4-dimethoxybenzyl bromide at a reaction temperature between about -10°C and about 60°C, preferably at room temperature to obtain N-
25 protected compound of formula VI. The deprotection reaction is carried out as described above.


For the synthesis of compounds of formula Ia or Ib wherein R^2 , R^3 , X and Z are as defined for compounds of formula I and R^1 is hydrogen, compound of formula III wherein R^1 is a N-protecting group known from the literature such as 4-methoxybenzyl,
30 benzyl or 3,4-dimethoxybenzyl (commercially available or synthesized according to methods known from the literature) is reacted according reaction steps 1-6 (and optionally 7) and finally deprotected under conditions known from the literature. The protecting reaction is for example carried out in polar aprotic solvents like dimethylsulfoxide (DMSO), dimethylacetamide or N,N-dimethylformamide (DMF), preferably
35 dimethylformamide in the presence of a base such as lithium hydride or sodium hydride, preferably sodium hydride at a reaction temperature between about -10°C and about 60°C,

preferably at room temperature. The reaction system is then treated with an appropriate protecting agent such as 4-methoxybenzyl chloride, 4-methoxybenzyl bromide, benzyl chloride, benzyl bromide, 3,4-dimethoxybenzyl chloride or 3,4-dimethoxybenzyl bromide at a reaction temperature between about -10°C and about 60°C, preferably at room temperature to obtain N-protected compound of formula VI.

The deprotecting reaction is carried out for example under hydrogen in the presence of a catalyst such as palladium on charcoal or platinum on charcoal (as described above). The corresponding compound of formula Ia or Ib wherein R², R³, X and Z are as defined for compounds of formula I and R¹ is hydrogen is obtained.

Optionally, compounds of formula Ia or Ib wherein R³, X and Z are as defined for compounds of formula I, R¹ or R² is hydrogen and the other one of R¹ or R² is as defined for compounds of formula I, can also be converted to the corresponding compound of formula Ia or Ib wherein R¹, R², R³, X and Z are as defined for compounds of formula I. The reaction is conveniently carried out with an electophilic agent R¹-L or R²-L, wherein R¹ and R² is as defined for formula I but not hydrogen and L is a leaving group such as bromide, chloride, iodide, tosylate, triflate or any other according to methods known from textbooks on organic chemistry e.g. from J. March (1992), "Advanced Organic Chemistry: Reactions, Mechanisms and Structure", 4th ed. John Wiley and Sons). The reaction is carried out as described in the literature, for example in the presence of a base such as sodium hydride, lithium hydride, potassium carbonate or triethylamine in an appropriate organic solvent such as tetrahydrofuran (THF) or polar aprotic solvents like dimethylsulfoxide (DMSO), N,N-dimethylacetamide or N,N-dimethylformamide (DMF), preferably DMF or THF at a reaction temperature between about -10°C and about 60°C, preferably at room temperature.

Reaction scheme 2:



wherein R^1 , R^2 , R^3 and X are as defined for compounds of formula I and R^4 is alkyl, aryl, optionally substituted phenyl or optionally substituted heterocyclyl as defined for compounds of formula I or C_{0-3} -alkyl substituted with C_{1-4} -alkoxy, C_{1-4} -alkyl-carbonyl-oxy- or heterocyclyl-oxy or R^4 together with the methyl group forms a optionally substituted C_{2-4} -alkenyl as defined for compounds of formula I.

In reaction scheme 2, the first reaction step is carried out in that the aldehydes of formula VIII are derivatised with a Grignard reagent of formula R^4MgHal wherein R^4 is alkyl, aryl, optionally substituted phenyl or optionally substituted heterocyclyl as defined for compounds of formula I or C_{0-3} -alkyl substituted with C_{1-4} -alkoxy, C_{1-4} -alkyl-carbonyl-oxy- or heterocyclyl-oxy or R^4 together with the methyl group forms a optionally substituted C_{2-4} -alkenyl as defined for compounds of formula I, and Hal represents chlorine, bromine or iodine, preferably chlorine (commercially available or synthesised according to textbooks on organic chemistry e.g. from J. March (1992), "Advanced Organic Chemistry: Reactions, Mechanisms and Structure", 4th ed. John Wiley and Sons) to obtain the corresponding alcohol derivative of formula Ic. The derivatisation reaction is conveniently carried out in an inert solvent for example ethers such as tetrahydrofuran, diethyl ether, dibutyl ether, dioxane, diglyme or a mixture of the mentioned solvents, preferably tetrahydrofuran at a reaction temperature between about -10°C and about 60°C, preferably at room temperature. In general, the derivatisation reaction can also be carried out as described in textbooks about organic chemistry e.g. from J. March (1992), "Advanced Organic Chemistry: Reactions, Mechanisms, and Structure", 4th ed. John Wiley & Sons.

In the second step of the reaction the methyl hydroxy group of alcohol derivative of formula Ic is converted via a deoxygenation reaction to the corresponding methylene group to obtain compound of formula Id. The reaction is conveniently carried out in the presence of trialkylsilane such as trimethylsilane, triethylsilane or tripropylsilane, preferably triethylsilane, dissolved in mineral acids such as trifluoroacetic acid (TFA) or in Lewis acids such as $SnCl_4$ (described in D. L. Comins et al., Tet. Lett., 1986, 27, 1869), preferably with trifluoroacetic acid at a reaction temperature from 0°C to 80°C, preferably at a reaction temperature between about 5°C and about 50°C. Alternatively, the reduction reaction can also be carried in the presence P2I4 as described in EP 0627423.


For the synthesis of compounds of formula Ic or Id wherein the substituents are as defined above and R^2 is hydrogen, the corresponding compound of formula VI is N-

protected as described for reaction scheme 1, further reacted according to reaction steps 4-5 of reaction scheme 1, then reacted according to reaction scheme 2 and finally deprotected as described for reaction scheme 1.

For the synthesis of compounds of formula Ic or Id wherein the substituents are as defined above and R¹ is hydrogen, compound of formula VIII wherein R¹ is a N-protecting group such as 4-methoxybenzyl (prepared according to reaction scheme 1) is reacted according reaction scheme 2 and finally deprotected as described for reaction scheme 1.

Optionally, compounds of formula Ic or Id wherein the substituents are as defined above and R¹ or R² is hydrogen and the other one of R¹ or R² is as defined above, can also be converted to the corresponding compound of formula Ic or Id wherein R¹, R², R³, X and Z are as defined above. The reaction is carried out as described for reaction scheme 1.

Reaction scheme 3:



wherein R, R¹, R², R³, X and Z are as defined for compounds of formula I.


In reaction scheme 3, the first reaction step is carried out in that aldehyde of formula VIII is derivatised via a reductive amination reaction with a primary or secondary amine of formula RN(H)CH₂-Z wherein R and Z are as defined for compounds of formula I to the corresponding amine compounds of formula Ic. The amines of formula RN(H)CH₂-Z are commercially available or can be synthesized according to known methods from textbooks on organic chemistry e.g. from J. March (1992), "Advanced Organic Chemistry: Reactions, Mechanisms and Structure", 4th ed. John Wiley and Sons. The reductive amination reaction is known from textbooks about organic chemistry e.g. from J. March (1992), "Advanced Organic Chemistry: Reactions, Mechanisms, and Structure", 4th ed. John Wiley & Sons. The reaction may be carried out in an inert solvent such as hydrocarbons for

example cyclohexane, methyl cyclohexane, decaline, benzene, toluene, o-xylene, m-xylene or p-xylene, preferably toluene at a reaction temperature between about 60°C and about boiling temperature of the organic solvent. After the reaction system has been cooled down, the organic solvent is evaporated and the residue is dissolved in an alcoholic solvent or ether for example methanol, ethanol, propanol, butanol, octanol or cyclohexanol, tetrahydrofuran, diethyl ether, dibutyl ether, dioxane or diglyme, preferably methanol, ethanol or tetrahydrofuran or a mixture of the mentioned solvents. The pH is adjusted in the range from 4.0 to 6.0, preferably in the range from 4.5 to 5.5 with inorganic acids, e.g. with hydrohalic acids such as hydrochloric acid, hydrobromic acid, sulphuric acid, nitric acid or phosphoric acid, and the like or with organic acids, e.g. with acetic acid, formic acid, tartaric acid, succinic acid, fumaric acid, maleic acid, malic acid, salicylic acid, citric acid, methanesulphonic acid and p-toluene sulphonic acid and the like. Subsequently, the system was reacted with a suitable reducing agent such as cyanoborohydride, sodium borohydride, NaBH₄, BH₃-THF, BH₃-SMe₂, catecholborane or other known in the art, preferably sodium cyanoborohydride or NaBH(OAc)₃ at a reaction temperature between about -10°C and about 60°C, preferably at room temperature.

For the synthesis of compounds of formula Ie wherein the substituents are as defined above and R² is hydrogen, the corresponding compound of formula VI is N-protected with an appropriate protecting group as described for reaction scheme 1, further reacted according to reaction steps 4-5 of reaction scheme 1, then reacted according to reaction scheme 3 and finally deprotected as described for reaction scheme 1.

For the synthesis of compounds of formula Ie wherein the substituents are as defined above and R¹ is hydrogen, compound of formula VIII wherein R¹ is a N-protecting group such as 4-methoxybenzyl (prepared according to reaction scheme 1) is reacted according to reaction scheme 3 and finally deprotected as described for reaction scheme 1.

Optionally, compounds of formula Ie wherein the substituents are as defined for compounds of formula I and R¹ or R² is hydrogen and the other one of R¹ or R² is as defined for compounds of formula I, can also be converted to the corresponding compound of formula Ie wherein R¹, R², R³, X and Z are as defined for compounds of formula I. The reaction is carried out as described for reaction scheme 1.

Reaction scheme 4:

wherein R^1 , R^2 , R^3 and X are as defined for compounds of formula I and R^5 is hydrogen, alkyl, protected hydroxy- C_{0-2} -alkyl or C_{0-2} -alkyl substituted with optionally substituted phenyl or with optionally substituted heterocyclyl as defined for compounds of formula I or C_{0-2} -alkyl substituted with C_{1-4} -alkoxy, C_{1-4} -alkyl-carbonyl-oxy- or heterocyclyl-oxy.

In reaction scheme 4, the first reaction step is carried out in that the aldehyde function of compound of formula VIII reacted via a Wittig-Horner reaction with dialkyl phosphonate of formula $(EtO)_2P(=O)(CH_2)R^5$ wherein R^5 is as defined above (commercially available or synthesised according to known methods from textbooks on organic chemistry e.g. from J. March (1992), "Advanced Organic Chemistry: Reactions, Mechanisms and Structure", 4th ed. John Wiley and Sons), to obtain olefinic compound of formula If. The reaction is carried out similar to the method described in the literature, for example in the presence of a strong base such as $n\text{-BuLi}$ or preferably sodium hydride in an organic solvent for example anhydrous ethers such as diethyl ether, dibutyl ether, dioxane, preferably anhydrous tetrahydrofuran under inert atmosphere such as nitrogen or argon atmosphere at a reaction temperature from 0°C to 80°C , preferably at a reaction temperature between about 5°C and about 50°C . Optionally, olefinic compound of formula If can be obtained through other coupling reactions for example the Wittig reaction.

In the second step of the reaction, the olefinic group of compound of formula If is hydrogenated to the corresponding compound of formula Ig. The reaction is carried out similar to methods described in the literature, for example under hydrogen in the presence of a hydrogenation catalyst in an appropriate solvent at a reaction temperature from 0°C to 80°C , preferably at a reaction temperature between about 5°C and about 50°C . The hydrogen pressure can be between about 0atm and about 100atm, preferably between about 0atm and about 50atm and most preferred between about 0atm and about 20atm.

The hydrogenation catalyst used for this reaction can be one of the commonly known catalysts such as noble metals (e.g. Pt, Pd or Rh) on supporting materials such as activated carbon or Al_2O_3 , or generally as described in textbooks about organic chemistry e.g. J. March (1992), "Advanced Organic Chemistry: Reactions, Mechanisms, and Structure", 4th ed. John Wiley & Sons. Preferred hydrogenation catalysts are Pd on activated carbon or Raney-Nickel. Appropriate solvents for the hydrogenation reaction are organic solvents such as alcohols (e.g. methanol, ethanol, propanol, butanol, octanol or cyclohexanol), ethers (e.g. tetrahydrofuran, diethyl ether, dibutyl ether or dioxane), polar aprotic solvents such as dimethylsulfoxide (DMSO) or N,N-dimethylacetamide, esters (e.g. ethyl acetate), halogenated hydrocarbons (e.g. dichloromethane or trichloromethane), hydrocarbons (e.g. cyclohexane, methyl cyclohexane, decaline, benzene, toluene, o-xylene, m-xylene or p-xylene) or a mixtures of the mentioned solvents. Preferred solvents are ester, most preferred solvent is ethyl acetate.

For the synthesis of compounds of formula Ig wherein R^5 is unprotected hydroxy- C_{0-2} -alkyl, the protected hydroxy group of compound of formula Ig is deprotected according to methods known from the art for example as described in 'Protecting Groups in Organic Synthesis', second edition, Theodora W. Greene and Peter G. M. Wuts, Wiley-Interscience, 1991.

For the synthesis of compounds of formula Ig wherein the substituents are as defined above and R^2 is hydrogen, the corresponding compound of formula VI is N-protected with an appropriate protecting group as described for reaction scheme 1, further reacted according to reaction steps 4-5 of reaction scheme 1, then reacted according reaction scheme 4 and finally deprotected as described for reaction scheme 1.

For the synthesis of compounds of formula Ig wherein the substituents are as defined above and R^1 is hydrogen, compound of formula VIII wherein R^1 is a N-protecting group such as 4-methoxybenzyl (prepared according to reaction scheme 1) is reacted according to reaction scheme 4 and finally deprotected under conditions known from the literature for example under hydrogen in the presence of a catalyst such as palladium on charcoal or platinum on charcoal.

Optionally, compounds of formula Ig wherein the substituents are as defined above and R^1 or R^2 is hydrogen and the other one of R^1 or R^2 is as defined above, can also be converted to the corresponding compound of formula Ig wherein the substituents are as defined above. The reaction is carried out as described for reaction scheme 1.

Assay Method: HIV-1 reverse transcriptase assay: Inhibitor IC₅₀ determination.

HIV-1 RT assay was carried out in 96-well Millipore filtermat NOB50 plates using purified recombinant enzyme and a poly(rA)/oligo(dT)₁₆ template-primer in a total volume of 50 µL. The assay constituents were 50 mM Tris/HCl, 50 mM NaCl, 1 mM
5 EDTA, 6 mM MgCl₂, 5 µM dTTP, 0.1 µCi [³H] dTTP, 5 µg/ml poly (rA) pre annealed to 2.5 µg/ml oligo (dT)₁₆ and a range of inhibitor concentrations in a final concentration of 10% DMSO. Reactions were initiated by adding 5 nM HIV-1 RT and after incubation at 37°C for 30 min, they were stopped by the addition of 50 µl ice cold 20%TCA and allowed to precipitate at 4°C for 30 min. The precipitates were collected by applying vacuum to the
10 plate and sequentially washing with 2 x 200 µl of 10% TCA and 2 x 200 µl 70% ethanol. Finally the plates were dried and radioactivity counted in a Wallac Microbeta 1450 after the addition of 15 µl scintillation fluid per well. IC₅₀'s were calculated by plotting % inhibition versus log₁₀ inhibitor concentrations.

Antiviral assay method

15 Anti-HIV antiviral activity was assessed using an adaptation of the method of Pauwels et al. {Pauwels et al., 1988, J Virol Methods 20:309-321}. The method is based on the ability of compounds to protect HIV-infected T lymphoblastoid cells (MT4 cells) from cell-death mediated by the infection. The endpoint of the assay was calculated as the concentration of compound at which the cell viability of the culture was preserved by 50%
20 ('50% inhibitory concentration', IC₅₀). The cell viability of a culture was determined by the uptake of soluble, yellow 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT) and its reduction to a purple insoluble formazan salt. After solubilization, spectrophotometric methods were employed to measure the amount of formazan product.

MT4 cells were prepared to be in logarithmic-phase growth and a total of 2 x 10⁶
25 cells infected with the HXB2-strain of HIV at a multiplicity of 0.0001 infectious units of virus per cell in a total volume of between 200-500 microlitres. The cells were incubated with virus for one h at 37°C before removal of virus. The cells are then washed in 0.01 M phosphate buffered saline, pH 7.2 before being resuspended in culture medium for incubation in culture with serial dilutions of test compound. The culture medium used
30 was RPMI 1640 without phenol red, supplemented with penicillin, streptomycin, L-glutamine and 10% fetal calf serum (GM10).

Test compounds were prepared as 2 mM solutions in dimethyl sulfoxide (DMSO). Four replicate, serial 2-fold dilutions in GM10 were then prepared and 50 microlitres amounts placed in 96-well plates over a final nanomolar concentration range of 625 – 1.22.

Fifty microlitres GM10 and 3.5×10^4 infected cells were then added to each well. Control cultures containing no cells (blank), uninfected cells (100% viability; 4 replicates) and infected cells without compound (total virus-mediated cell death; 4 replicates) were also prepared. The cultures were then incubated at 37 °C in a humidified atmosphere of 5% CO₂ in air for 5 s.






A fresh solution of 5 mg/ml MTT was prepared in 0.01 M phosphate buffered saline, pH 7.2 and 20 microlitres added to each culture. The cultures were further incubated as before for 2 h. They were then mixed by pipetting up and down and 170 microlitres of Triton X-100 in acidified isopropanol (10% v/v Triton X-100 in 1:250 mixture of concentrated HCl in isopropanol). When the formazan deposit was fully solubilized by further mixing, the absorbance (OD) of the cultures was measured at 540nm and 690nm wavelength (690nm readings were used as blanks for artefacts between wells). The percent protection for each treated culture was then calculated from the equation:







% Protection =





$$\frac{(\text{OD drug-treated cultures}) - (\text{OD untreated virus control cultures})}{(\text{OD uninfected cultures}) - (\text{OD untreated virus control cultures})} \times 100\%$$

The IC₅₀ was then obtained from graph plots of percent protection versus log₁₀ drug concentration.

In the assay, compounds of formulas I range in activity from an HIV IC₅₀ of about 0.5 to about 5000 nM, with preferred compounds having a range of activity from about 0.5 to about 750 nM, more preferably about 0.5 to 300 nM, and most preferably about 0.5 to 50 nM.

Structure	RT IC ₅₀ [nM]	HIV IC ₅₀ [nM]
	90	7
	171	4
	520	32
	730	44
	2400	108

	1160	ND
	1905	ND
	3339	ND
	10000	ND
	10000	ND
	1919	ND

	649	ND
	313	7
	608	134
	115	5

ND: not determined

The imidazolone derivatives provided by the present invention can be used together with a therapeutically inert carrier as medicaments in the form of pharmaceutical preparations. The pharmaceutical preparations can be administered enterally, such as orally, in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions, or nasally, e.g. in the form of nasal sprays. They can also be administered rectally, e.g. in the form of suppositories, or parenterally, (e.g. intramuscularly, intravenously, or subcutaneously), for example, in the form of injection solutions.

For the manufacture of pharmaceutical preparations the imidazolone derivatives can be formulated with therapeutically inert, inorganic or organic carriers.

Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts can be used, for example, as such carriers for tablets, coated tablets, dragées and hard gelatine capsules.

Suitable carriers for soft gelatine capsules are, for example, vegetable oils, waxes, fats, semi-solid and liquid polyols and the like.

Suitable carriers for the manufacture of injection solutions are, for example, water, saline, alcohols, polyols, glycerine, vegetable oils and the like. Natural or hardened oils, waxes, fats, semi-liquid or liquid polyols and the like are suitable carriers for the manufacture of suppositories. The pharmaceutical preparations of the present invention may also be provided as sustained release formulations or other appropriate formulations.

The pharmaceutical preparations can also contain preservatives, solubilizers, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavourants, salts for adjustment of the osmotic pressure, buffers, masking agents or antioxidants.

The pharmaceutical preparations may also contain other therapeutically active agents such as those mentioned above.

The imidazolone derivatives provided by the invention in the treatment of an immune mediated condition or disease, a viral disease, a bacterial disease, a parasitic disease, an inflammatory disease, a hyperproliferative vascular disease, a tumor, or cancer.

The dosage can vary within wide limits and will, of course, be adjusted to the individual requirements in each particular case.

Dosage levels of between about 0.01 and about 100 mg/kg body weight per day in monotherapy and/or in combination therapy are commonly administered from about 1 to

- 50 -

5 times per day. A typical preparation will contain from about 5% to 95% active compound (w/w) . The daily dosage can be administered as a single dosage or in divided dosages.

The imidazolone derivatives provided by the present invention or the medicaments
5 thereof may be for use in monotherapy and/or combination therapy, i.e. the treatment
may be in conjunction with the administration of one or more additional therapeutically
active substance(s). When the treatment is combination therapy, such administration may
be concurrent or sequential with respect to that of the imidazolone derivatives of the
present invention. Thus, concurrent administration, as used herein, includes
10 administration of the agents in conjunction or combination, together, or before or after
each other.

It will be understood that references herein to treatment extend to prophylaxis as
well as to treatment of existing conditions. Treatment of a disease or condition, as used
herein, also includes preventing, inhibiting, regressing, reversing, alleviating or relieving
15 the disease or condition, or the clinical symptoms thereof. The term "subject" as used
herein refers to animals, including humans and other mammals.

In the following examples the abbreviations used have the following significations:

	MS	mass spectroscopy
	ES	electrospray
	EI	electron impact
5	NMR	nuclear magnetic resonance spectroscopy
	DMF	N,N-dimethylformamide
	DMSO	dimethylsulfoxide
	rt	room temperature
	min	minute(s)
10	h	hour(s)
	d	day(s)

All temperatures are given in degrees Celsius (°C).

The described NMR spectra were recorded on a Bruker DRX 400 MHz spectrometer
15 with the probe temperature set at 300 K.

The mass spectra indicated by "(M⁺; EI)", were recorded under electron impact conditions (EI), on a THERMOQUEST MAT95 S with a source temperature of 200°C. Other mass spectra were recorded under electrospray ionization spectra (ESI) conditions, on one of the following machines:

20 a) THERMOQUEST SSQ 7000 [Solvent 0.085% TFA in 90% Acetonitrile/water; flow rate 100 microliters/minute; capillary 250°C; spray voltage 5KV; sheath gas 80 psi], or

b) LC-MS system (liquid chromatograph coupled to mass spectrum)
THERMOQUEST TSQ 7000 ELECTROSPRAY or MICROMASS PLATFORM
ELECTROSPRAY [Solvent 0.1% TFA in water or 0.085% TFA in 90% acetonitrile/ water
25 or 0.085% TFA in acetonitrile].

With regard to the starting materials that are known compounds some of these may be purchased from commercial suppliers. Other starting materials that are known and their analogues can be prepared by methods well known in the art. Examples of compounds available from commercial suppliers, and citations to the synthesis of other
5 compounds and their analogues are provided in the following:

Compounds, whenever prepared by the processes of the present invention are also an object of the present invention.

The following examples illustrate the present invention:

10

Example 1

4-(3,5-Dichlorophenylthio)-5-(hydroxymethyl)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone

A mixture of 117mg of 5-(3,5-dichlorophenylthio)-1-isopropyl-3-methyl-1,3-dihydro-2-imidazolone-4-carboxaldehyde and 33mg of sodium borohydride in 4ml of
15 tetrahydrofuran and 1ml of ethanol was stirred at rt for 3 h. The reaction mixture was reduced in volume and the residue was purified by flash chromatography on silica gel using ethyl acetate/hexane (1:1 then 2:1 then 100% ethyl acetate) for the elution to give 113mg of white crystals. Mass spectrum (ES) m/z 347 [M+H]⁺.

20 The starting material was prepared as follows:

Intermediate 1A

5-(3,5-dichlorophenylthio)-1-isopropyl-3-methyl-1,3-dihydro-2-imidazolone-4-carboxaldehyde

25 To a solution of 4.16g of 5-chloro-1-isopropyl-3-methyl-1,3-dihydro-2-imidazolone-4-carboxaldehyde in 50ml of dichloromethane was added 4.4g of 3,5-dichlorothiophenol and 6.8ml of triethylamine. The mixture was stirred at rt under nitrogen for 18 h. The reaction mixture was concentrated and partitioned between ethyl acetate and 2N hydrochloric acid. The organic layer was washed with a further aliquot of

2N hydrochloric acid and brine then dried, filtered and evaporated to dryness. The residue crystallised on standing to provide a solid, which was triturated with diethyl ether to give 5g of a white solid. Mass spectrum (ES) m/z 344 $[M+H]^+$.

5

Intermediate 1B

5-Chloro-1-isopropyl-3-methyl-1,3-dihydro-2-imidazolone-4-carboxaldehyde

A solution of 6.5g of 5-dimethylaminomethylene-3-isopropyl-1-methyl-imidazolidine-2,4-dione in 30ml of phosphorus oxychloride and 5ml of dimethylformamide was heated to 100°C for 2 h. After this time, the reaction mixture was cooled to rt, poured into ice water and extracted with ethyl acetate. The organic extract was dried, filtered and evaporated to give 4.16g as yellow needles. Mass spectrum (EI) m/z 202 M^+ .

10

Intermediate 1C

5-Dimethylaminomethylene-3-isopropyl-1-methyl-imidazolidine-2,4-dione

A mixture of 13.4g of 3-isopropyl-1-methyl-imidazolidine-2,4-dione and 13.6ml dimethylformamide dimethylacetal were heated at 100°C over the weekend. The reaction mixture was cooled to rt and purified by flash chromatography to give 9.9g of a golden oil. Mass spectrum (EI) m/z 211 M^+ .

20

Intermediate 1D

3-Isopropyl-1-methylimidazolidine-2,4-dione

A mixture of 14.9g of (3-isopropyl-1-methylureido)acetic acid in 50ml of water and 50ml of concentrated hydrochloric acid was refluxed for 2 h. On cooling the mixture was diluted with water and extracted with ethyl acetate to give 13.4g of a colourless oil. Mass spectrum (EI) m/z 156 M^+ .

25

Intermediate 1E(3-Isopropyl-1-methylureido)acetic acid

To a solution of 4g of sodium hydroxide in 75ml of water was added 10g of sarcosine portionwise. The solution was cooled and treated with 11.5ml of isopropyl isocyanate.

- 5 After 2 h, the reaction mixture was washed with diethyl ether and the aqueous phase acidified with concentrated HCl to <pH 2. The product was extracted twice with ethyl acetate to give 10.45g of a white solid which was taken on to the next step.

A further extraction of the aqueous phase after standing gave 5.5g of a colourless oil the next product.

10

Example 25-(Benzyloxymethyl)-4-(3,5-dichlorophenylthio)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone

- To 50mg of 4-(3,5-dichlorophenylthio)-5-(hydroxymethyl)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone in 1ml of dimethylformamide was added 7mg of sodium
15 hydride (60% dispersion in oil). The reaction mixture was stirred at rt for 30 min until hydrogen gas evolution had ceased. The solution was then treated with 20µL of benzyl bromide and stirring continued for 1 h. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was separated, dried and evaporated to dryness
20 to give 190mg of an oily residue. This residue was purified by flash chromatography on silica gel using ethyl acetate/hexane (1:1) for the elution to give 55mg of a colourless oil. Mass spectrum (ESI) m/z 437 [M+H]⁺.

Example 3

- 25 5-(Benzyloxymethyl)-4-(3,5-dichlorophenoxy)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone

To 115mg of 4-(3,5-dichlorophenoxy)-5-hydroxymethyl-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone in 5ml of dimethylformamide was added 15mg of sodium hydride (60% dispersion in oil). The reaction mixture was stirred at rt for 30 min until hydrogen

gas evolution had ceased. The solution was then treated with 45 μ L of benzyl bromide and stirring continued for 1 h. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was separated, dried and evaporated to dryness to give 190mg of an oily residue. This residue was purified by flash chromatography on silica gel using ethyl acetate/hexane (1:1) for the elution to give 41mg of a colourless oil. Mass spectrum (ESI) m/z 421 [M+H]⁺.

The starting material was prepared as follows:

10

Example 3A

4-(3,5-Dichlorophenoxy)-5-hydroxymethyl-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone

A mixture of 100mg of 5-(3,5-dichlorophenoxy)-1-isopropyl-3-methyl-1,3-dihydro-2-imidazolone-4-carboxaldehyde and 13mg of sodium borohydride in 5ml of methanol was stirred at rt for 2 h. The reaction mixture was reduced in volume then dissolved in ethyl acetate. The solution was washed with water, dried and evaporated to dryness to give 115mg of colourless needles. Mass spectrum (ESI) m/z 331 [M+H]⁺.

Example 3B

5-(3,5-Dichlorophenoxy)-1-isopropyl-3-methyl-2,3-dihydro-2-imidazolone-4-carboxaldehyde

To a solution of 163mg of 3,5-dichlorophenol in 3ml of dimethylformamide was added portionwise 26mg of sodium hydride (as a 60% dispersion in mineral oil). The reaction mixture was stirred until evolution of hydrogen gas has ceased then was treated with 200mg of 5-chloro-1-isopropyl-3-methyl-1,3-dihydro-2-imidazolone-4-carboxaldehyde. The mixture was stirred overnight at rt. The reaction mixture was diluted with ethyl acetate and washed with dilute sodium hydroxide solution. The organic phase was dried, filtered and evaporated to give 341mg of a buff solid. Mass spectrum (ESI) m/z 437, 438 [M+H]⁺.

Example 44-(3,5-Dichlorophenylthio)-5-(hydroxymethyl)-3-isopropyl-1-(4-methoxybenzyl)-1,3-dihydro-2-imidazolone

- 5 A mixture of 50mg of 5-(3,5-dichlorophenylthio)-1-isopropyl-3-(4-methoxybenzyl)-1,3-dihydro-2-imidazolone-4-carboxaldehyde and 5mg of sodium borohydride in 5ml of methanol was stirred at rt for 2 h. The reaction mixture was reduced in volume and dissolved in ethyl acetate, washed with water, dried and evaporated to give 50mg of a white solid. Mass spectrum (ESI) m/z 453 [M+H]⁺.

10

The starting material was prepared as follows:

Intermediate 4A

- 15 5-(3,5-Dichlorophenylthio)-1-isopropyl-3-(4-methoxybenzyl)-1,3-dihydro-2-imidazolone-4-carboxaldehyde

- 20 To a solution of 1.99g of 5-chloro-1-isopropyl-3-(4-methoxybenzyl)-1,3-dihydro-2-imidazolone-4-carbaldehyde in 25ml of dichloromethane was added 1.2g of 3,5-dichlorothiophenol and 2ml of triethylamine. The mixture was stirred at rt under nitrogen for 18 h. The reaction mixture was concentrated and partitioned between ethyl acetate and 2N hydrochloric acid. The organic layer was washed with a further aliquot of 2N hydrochloric acid and brine then dried, filtered and evaporated to dryness. The residue crystallised on standing to provide a solid, which was triturated with diethyl ether to give 1.4g of a white solid. Mass spectrum (ESI) m/z 451 [M+H]⁺.

Intermediate 4B5-Chloro-1-isopropyl-3-(4-methoxybenzyl)-1,3-dihydro-2-imidazolone-4-carboxaldehyde

A mixture of 1.99g of 5-dimethylaminomethylene-3-isopropyl-1-(4-methoxybenzyl)imidazolidine-2,4-dione, 10ml of phosphorus oxychloride and 2ml of dimethylformamide was heated at 100°C for 2 h. The reaction mixture was cooled to rt, poured onto ice water and extracted with ethyl acetate. After evaporation the crude product was purified by flash chromatography to give 2.2g of an oil. Mass spectrum (ESI) m/z 309 [M+H]⁺.

10

Intermediate 4C5-Dimethylaminomethylene-3-isopropyl-1-(4-methoxybenzyl)-imidazolidine-2,4-dione

To a solution of 100mg of 5-dimethylaminomethylene-3-isopropylimidazolidine-2,4-dione in 2ml of dimethylformamide was added 425mg of sodium hydride (as a 60% dispersion in mineral oil). The mixture was stirred until evolution of hydrogen gas had ceased. The solution was then treated with 1.5ml of 4-methoxybenzyl chloride and stirring continued overnight at rt. The mixture was partitioned between ethyl acetate and water and the organic layer dried, filtered and concentrated. The residue was purified by flash chromatography eluting with 1:1 ethyl acetate: hexane to give 85mg of a colourless oil. ¹H NMR (400MHz, deuteriochloroform): δ = 1.40 (6H, d, J = 7 Hz), 3.02 (6H, s), 3.72 (3H, s), 4.38 (1H, septet, J = 7 Hz), 4.60 (2H, s), 5.96 (1H, s), 6.80 (2H, d, J = 8 Hz), 7.05 (2H, d, J = 8 Hz).

15
20Intermediate 4D5-Dimethylaminomethylene-3-isopropylimidazolidine-2,4-dione

25

A mixture of 5g of 3-isopropylimidazolidine-2,4-dione and 5ml dimethylformamide dimethylacetal were heated at 100°C for 3 s. The reaction mixture was cooled to rt and purified by flash chromatography to give 2g of an orange solid. ¹H NMR (400MHz,

deuteriochloroform): δ = 1.42 (6H, d, J = 7 Hz), 3.11 (6H, s), 4.40 (1H, septet, J = 7 Hz), 6.64 (1H, s), 9.55 (1H, br s).

Intermediate 4E

5 3-Isopropylimidazolidine-2,4-dione

To a solution of 3.6g of sodium hydroxide in 75ml of water was added 7.5g of glycine. The solution was cooled to 5°C and treated with 9.8ml of isopropylisocyanate. After 5 h the reaction mixture was neutralised with 50ml concentrated hydrochloric acid added dropwise to the slurry. This mixture was refluxed for 2 h then cooled to rt. The
10 product was extracted with ethyl acetate to give 10.9g of a colourless oil. ¹H NMR (400MHz, deuteriochloroform): δ = 1.45 (6H, d, J = 7 Hz), 3.92 (2H, s), 4.35 (1H, septet, J = 7 Hz), 6.39 (1H, br s).

Example 5

15 4-(3,5-Dichlorophenylthio)-5-(2-hydroxybenzyl)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone







To a solution of 100mg of 4-(3,5-dichlorophenylthio)-5-(hydroxymethyl)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone, 30mg of phenol and 100mg of
20 triphenylphosphine in 5mL of tetrahydrofuran at 0°C was added 70μL of diisopropylazodicarboxylate in 5mL of tetrahydrofuran over 20 min. The reaction was stirred overnight warming to room temperature. The mixture was then diluted with ethyl acetate and washed with 0.5N hydrochloric acid solution then 0.5N sodium hydroxide solution, dried, concentrated and purified by flash chromatography eluting with 3:1
25 hexane: ethyl acetate to give 16mg of a colourless oil. Mass spectrum (ES) m/z 423 [M+H]⁺.

Examples 6-9 and 33-39

The compounds shown in table 2 were prepared in a manner analogous to that described for examples 1 to 5:

Table 2

Ex.	Structure	Name	MS (ES) (M + H)+
6		4-(3,5-Dichlorophenylthio)-5-hydroxymethyl-1-methyl-3-phenyl-1,3-dihydro-2-imidazolone	382
7		5-Benzyloxymethyl-4-(3,5-dichlorophenylthio)-1-methyl-3-phenyl-1,3-dihydro-2-imidazolone	471
8		4-(3,5-Dichlorophenylthio)-3-isopropyl-1-methyl-5-(pyridin-4-ylmethoxymethyl)-1,3-dihydro-2-imidazolone	438
9		3-Benzyl-4-(3,5-dichlorophenylthio)-5-hydroxymethyl-1-methyl-1,3-dihydro-2-imidazolone	395
33		4-(3,5-Dichloro-phenylsulfanyl)-3-isopropyl-5-methoxymethyl-1-methyl-1,3-dihydro-imidazol-2-one	362

34		4-Hydroxymethyl-3-methyl-1-phenyl-5-(pyridin-4-ylsulfanyl)-1,3-dihydro-imidazol-2-one	314
35		1-Benzyl-4-hydroxymethyl-3-methyl-5-(pyridin-4-ylsulfanyl)-1,3-dihydro-imidazol-2-one; hydrochloric acid	364
36		4-(3,5-Dichloro-phenylsulfanyl)-3-isopropyl-1-methyl-5-(pyridin-3-ylmethoxymethyl)-1,3-dihydro-imidazol-2-one	439
37		4-(4-Benzyloxy-benzyloxymethyl)-5-(3,5-dichloro-phenylsulfanyl)-1-isopropyl-3-methyl-1,3-dihydro-imidazol-2-one	544
38		Acetic acid 5-(3,5-dichloro-phenylsulfanyl)-1-isopropyl-3-methyl-2-oxo-2,3-dihydro-1H-imidazol-4-ylmethyl ester	390
39		4-(3,5-Dichloro-phenylsulfanyl)-3-isopropyl-1-methyl-5-(pyridin-4-ylloxymethyl)-1,3-dihydro-imidazol-2-one	425

Example 104-(3,5-Dichlorophenylthio)-5-(alpha(RS)-hydroxybenzyl)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone

5 A mixture of 345mg of 5-(3,5-dichlorophenoxy)-1-isopropyl-3-methyl-1,3-dihydro-2-imidazolone-4-carboxaldehyde in 5ml of tetrahydrofuran was added to 0.4ml of a 3M solution of phenyl magnesium bromide also in 5ml of tetrahydrofuran under nitrogen. The reaction was stirred at rt for 2 h then quenched with saturated ammonium chloride solution and extracted with ethyl acetate. The organic phase was washed with brine, dried and evaporated to give 490mg of a solid foam. Mass spectrum (ESI) m/z 423 [M+H]⁺.

10

Example 114-(3,5-Dichlorophenylthio)-3-isopropyl-1-methyl-5-benzyl-1,3-dihydro-2-imidazolone




15 To a mixture of 250mg of 4-(3,5-dichlorophenoxy)-5-(alpha(RS)-hydroxybenzyl)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone in 3ml of trifluoroacetic acid was added 90μL of triethylsilane and stirred under nitrogen for 0.5 h. The reaction mixture was evaporated then the residue was purified by flash chromatography to give 245mg of an amber oil. Mass spectrum (ES) m/z 407 [M+H]⁺.

Examples 12 and 40-48

20 The compounds shown in table 3 were prepared in a manner analogous to that described for examples 10 to 11:

Table 3

Ex.	Structure	Name	MS (ES) (M + H)+
12		3-Benzyl-4-(3,5-dichlorophenylthio)-5-(1-hydroxyethyl)-1-methyl-1,3-dihydro-2-imidazolone	409
40		3-(5-Benzyl-3-isopropyl-1-methyl-2-oxo-2,3-dihydro-1H-imidazol-4-ylsulfanyl)-benzonitrile	364
41		4-Benzyl-5-isobutylsulfanyl-1-isopropyl-3-methyl-1,3-dihydro-imidazol-2-one	319
42		4-Benzyl-1-isopropyl-5-isopropylsulfanyl-3-methyl-1,3-dihydro-imidazol-2-one	305
43		4-Benzyl-1-isopropyl-3-methyl-5-methylsulfanyl-1,3-dihydro-imidazol-2-one	277
44		4-(3,5-Dichloro-phenylsulfanyl)-5-isobutyl-3-isopropyl-1-methyl-1,3-dihydro-imidazol-2-one	374
45		4-Benzyl-5-(3,5-dichloro-phenylsulfanyl)-3-ethyl-1-isopropyl-1,3-dihydro-imidazol-2-one	422

46		1-Allyl-5-benzyl-4-(3,5-dichloro-phenylsulfanyl)-3-isopropyl-1,3-dihydro-imidazol-2-one	434
47		4-(3,5-Dichloro-phenylsulfanyl)-1-ethyl-3-isopropyl-5-pyridin-4-ylmethyl-1,3-dihydro-imidazol-2-one	423
48		1-Allyl-4-(3,5-dichloro-phenylsulfanyl)-3-isopropyl-5-pyridin-4-ylmethyl-1,3-dihydro-imidazol-2-one	435

Example 13

5 4-(3,5-Dichlorophenylthio)-3-isopropyl-1-methyl-5-[(4-pyridyl)methylamino]methyl]-1,3-dihydro-2-imidazolone

A mixture of 100mg of 4-(3,5-dichlorophenylthio)-1-isopropyl-3-methyl-1,3-dihydro-2-imidazolone-4-carboxaldehyde and 34μL of 4-pyridylmethylamine in 10ml of toluene was azeotroped in a Dean-Stark apparatus for 2 h. The reaction was cooled to rt, evaporated and re-dissolved in 10ml of methanol. Bromocresol green was added and the pH adjusted to approximately 5 using 4N hydrochloric acid/dioxane. The solution was treated portionwise with 30mg of sodium cyanoborohydride and the pH adjusted back to 5 using 4N hydrochloric acid/dioxane. This was repeated until the addition was complete. The reaction was evaporated and the residue was purified by flash chromatography on silica gel using ethyl acetate/hexane (2:1) followed by methanol/dichloromethane (1:19) for the elution to give 72mg of a colourless oil. Mass spectrum (ES) m/z 437, 438 [M+H]⁺.

Example 144-(3,5-Dichlorophenylthio)-3-isopropyl-1-(4-methoxybenzyl)-5-[[4-pyridyl)methylamino]methyl]-1,3-dihydro-2-imidazolone






A mixture of 1g of 5-(3,5-dichlorophenylthio)-1-isopropyl-3-(4-methoxybenzyl)-
5 dihydro-2-imidazolone-4-carboxaldehyde and 0.25ml of 4-pyridylmethylamine in 20ml of
toluene was refluxed in a Dean-Stark apparatus for 2 h. The reaction was cooled to rt,
evaporated and re-dissolved in 10ml of methanol. Bromocresol green was added and the
pH adjusted to approximately 5 using 4N hydrochloric acid/dioxane. The solution was
treated portionwise with 155mg of sodium cyanoborohydride and the pH adjusted back to
10 5 using 4N hydrochloric acid/dioxane. This was repeated until the addition was complete.
The reaction was evaporated and the residue was purified by flash chromatography on
silica gel eluting with ethyl acetate/hexane (2:1) followed by methanol/dichloromethane
(1:19) to give 1.07g of a golden solid. Mass spectrum (ES) m/z 543 [M+H]⁺.






15

Examples 15-31



The compounds shown in table 4 were prepared in a manner analogous to that
described for examples 13 to 14:

Table 4

Ex.	Structure	Name	MS (ES) (M + H) ⁺
15		5-(Benzylaminomethyl)-4-(3,5-dichlorophenylthio)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone	477 ((M+MeCN+H) ⁺ ; ESI)
16		5-(Benzylaminomethyl)-3-isopropyl-1-methyl-4-phenylthio-1,3-dihydro-2-imidazolone	367
17		3-Isopropyl-1-methyl-4-phenylthio-5-[[pyridin-4-ylmethyl]amino]methyl-1,3-dihydro-2-imidazolone	369
18		5-(Benzylaminomethyl)-4-(3,5-dimethylphenylthio)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone	396
19		4-Cyclohexylthio-3-isopropyl-1-methyl-5-[[pyridin-4-ylmethyl]amino]methyl-1,3-dihydro-2-imidazolone	375

20		4-(3,5-Dimethylphenylthio)-3-isopropyl-1-methyl-5-(((pyridin-4-ylmethyl)amino)methyl)-1,3-dihydro-2-imidazolone	397
21		4-Benzylthio-3-isopropyl-1-methyl-5-(((pyridin-4-ylmethyl)amino)methyl)-1,3-dihydro-2-imidazolone	383
22		4-(3-Chlorophenylthio)-3-isopropyl-1-methyl-5-(((pyridin-4-ylmethyl)amino)methyl)-1,3-dihydro-2-imidazolone	403
23		3-Isopropyl-4-isopropylthio-1-methyl-5-(((pyridin-4-ylmethyl)amino)methyl)-1,3-dihydro-2-imidazolone	355
24		4-Isobutylthio-3-isopropyl-1-methyl-5-(((pyridin-4-ylmethyl)amino)methyl)-1,3-dihydro-2-imidazolone	349

25		3-Benzyl-4-(3,5-dichlorophenylthio)-1-methyl-5-[[[(pyridin-4-ylmethyl)amino]methyl]-1,3-dihydro-2-imidazolone	485
26		1,3-Dibenzyl-4-(3,5-dichlorophenylthio)-5-[[[(pyridin-4-ylmethyl)amino]methyl]-1,3-dihydro-2-imidazolone	561
27		1-Benzyl-4-(3,5-dichlorophenylthio)-3-isopropyl-5-[[[(pyridin-4-ylmethyl)amino]methyl]-1,3-dihydro-2-imidazolone	513
28		1-Benzyl-4-(3,5-dichlorophenylthio)-3-propyl-5-[[[(pyridin-4-ylmethyl)amino]methyl]-1,3-dihydro-2-imidazolone	513
29		4-(3,5-Dichlorophenoxy)-3-isopropyl-1-methyl-5-[[[(pyridin-4-ylmethyl)amino]methyl]-1,3-dihydro-2-imidazolone	421

30		4-sec-Butylthio-3-isopropyl-1-methyl-5- {[(pyridin-4-ylmethyl)amino]methyl}-1,3- dihydro-2-imidazolone	349
31		1-Isopropyl-5-(3-methoxyphenylthio)-3- methyl-4-{[(pyridin-4- ylmethyl)amino]methyl}-1,3-dihydro-2- imidazolone	399

Example 32

5 4-(3,5-Dichlorophenylthio)-3-isopropyl-1-methyl-5-(2-phenylethyl)-1,3-dihydro -2-imidazolone

A solution of 50mg of 4-(3,5-dichlorophenylthio)-5-styryl-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone was hydrogenated over a catalytic amount of 5% Pd on carbon for 4 h. The reaction mixture was filtered and evaporated then purified by flash chromatography eluting with 3:1 hexane/ethyl acetate to give 5mg of a colourless oil. Mass spectrum (ES) m/z 421 [M+H]⁺.

10

The starting material was prepared as follows:

Intermediate 32A

4-(3,5-Dichlorophenylthio)-5-styryl-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone

15 Polymer supported triphenylphosphine (1g, 3mmol) and 713μL of benzyl bromide were refluxed in 20ml of toluene for 3 h. The mixture was allowed to cool to rt and the toluene decanted. The resulting salt was stirred in 10ml of tetrahydrofuran at 0°C and treated with potassium tert.-butoxide. This mixture was stirred for 30 min at 0°C then

345mg of 5-(3,5-dichlorophenylthio)-2,3-dihydro-1-isopropyl-3-methyl-2-oxo-1H-imidazole-4-carboxaldehyde was added. The mixture was stirred at rt for 3 s then filtered and evaporated. The residue was purified by flash chromatography eluting with 3:2 hexane/ethyl acetate to give 140mg of a colourless oil. Mass spectrum (ES) m/z 419

5 $[M+H]^+$.

Examples 49-52

The compounds shown in table 5 were prepared in a manner analogous to that described for examples 13 to 14:

Table 5

Ex.	Structure	Name	MS (ES) (M + H)+
49		[E]-3-[5-(3,5-Dichloro-phenylsulfanyl)-1-isopropyl-3-methyl-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-acrylic acid ethyl ester	416
50		3-[5-(3,5-Dichloro-phenylsulfanyl)-1-isopropyl-3-methyl-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-acrylonitrile	369
51		3-[5-(3,5-Dichloro-phenylsulfanyl)-1-isopropyl-3-methyl-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-acrylamide	387
52		1-Benzyl-5-(3,5-dichloro-phenylsulfanyl)-3-methyl-4-vinyl-1,3-dihydro-imidazol-2-one	392

- 70 -

Example 534-(3,5-Dichloro-phenylsulfanyl)-3-isopropyl-1-methyl-5-pyridin-4-ylmethyl-1,3-dihydro-imidazol-2-one


A solution of 160mg of phosphorus tetraiodide in 15mL of toluene was heated in the
5 dark at 80°C for 20 min. To this solution was added dropwise a solution of 150mg of 4-
(3,5-dichloro-phenylsulfanyl)-3-isopropyl-1-methyl-5-[1-hydroxy-1-pyridin-4-ylmethyl]-
1,3-dihydro-imidazol-2-one in 5mL of toluene. The mixture was then heated at 80°C for 1
h then allowed to cool to room temperature. Then 8mL of 10% aqueous sodium bisulphite
10 solution was added and the biphasic mixture stirred for 1 h. The aqueous phase was
extracted three times with 30mL of ethyl acetate and combined extracts were washed with
brine, dried, filtered and evaporated to give a yellow residue which was purified by flash
chromatography on silica gel using ethyl acetate/methanol (19:1) for the elution to give
50mg as a pale yellow oil. Mass spectrum (ES) m/z 408 $[M+H]^+$.

The starting material 4-(3,5-dichloro-phenylsulfanyl)-3-isopropyl-1-methyl-5-[1-
15 hydroxy-1-pyridin-4-ylmethyl]-1,3-dihydro-imidazol-2-one was prepared as follows:

560mg of 4-Bromopyridine.hydrobromide was treated with 15mL of 5% aqueous
sodium hydrogen carbonate and extracted three times with 20mL of diethyl ether and
combined extracts were washed with brine, dried, filtered and evaporated to give a
colourless oil which was dissolved in 3mL of tetrahydrofuran. To this solution, under
20 nitrogen at room temperature, was added 1.45mL of a 3.0M solution of isopropyl
magnesium chloride in diethyl ether. The reaction mixture was stirred at room
temperature for 1.5 h then a solution of 1.0g of 5-(3,5-dichlorophenylthio)-1-isopropyl-3-
methyl-1,3-dihydro-2-imidazolone-4-carboxaldehyde in 10mL of tetrahydrofuran was
added. The reaction mixture was stirred at room temperature overnight then 20mL of
25 water added. The aqueous phase was extracted three times with 10mL of dichloromethane
and combined extracts were washed with brine, dried, filtered and evaporated to give a
yellow oil which was purified by flash chromatography on silica gel using ethyl
acetate/hexane (2:1 then 4:1) for the elution to give 250mg of a pale yellow solid. Mass
spectrum (ES) m/z 424 $[M+H]^+$.

Claims

1. Compounds of formula I



wherein

5 R^1 is hydrogen, alkyl, cycloalkyl, aryl or alkyl substituted with optionally substituted phenyl;

R^2 is hydrogen, alkenyl, alkyl or alkyl substituted with optionally substituted phenyl;

R^3 is alkyl, cycloalkyl, optionally substituted phenyl, optionally substituted benzyl or heterocyclyl;

10 A is alkyl, hydroxy-alkyl, alkenyl, substituted alkenyl, aryl-CH(OH)-, substituted alkyl, or

A signifies a group of formula $Z-CH_2-Y-CH_2-$,

wherein Y represents O or NR wherein R is hydrogen or alkyl and

Z represents heterocyclyl or optionally substituted aryl;

15 X represents S or O;

with the provisos that

(i) only one of R^1 and R^2 is hydrogen;

(ii) when X is O, then R^1 cannot be aryl;

as well as hydrolyzable esters or ethers of compounds of formula I and pharmaceutically
20 acceptable salts thereof.

2. Compounds as claimed in claim 1 wherein

- 72 -

R¹ is hydrogen, C₁₋₁₂-alkyl, C₃₋₈-cycloalkyl, aryl or C₁₋₄-alkyl substituted with phenyl optionally substituted with 1-5 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine;

5 R² is hydrogen, C₂₋₄-alkenyl, C₁₋₁₂-alkyl or C₁₋₄-alkyl substituted with phenyl optionally substituted with 1-5 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine;

R³ is C₁₋₁₂-alkyl, C₃₋₈-cycloalkyl, phenyl, benzyl or heterocyclyl,
wherein a phenyl ring may be substituted with 1-5 substituents selected
from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine, bromine and
10 cyano;

A is C₁₋₁₂-alkyl, hydroxy-C₁₋₄-alkyl, C₂₋₄-alkenyl, substituted C₂₋₄-alkenyl, aryl-CH(OH)-, C₁₋₄-alkyl substituted with C₁₋₄-alkoxy, C₁₋₄-alkyl-carbonyl-oxy-, heterocyclyl-oxy or A is C₁₋₄-alkyl substituted with phenyl optionally substituted
with 1-5 substituents or with heterocyclyl optionally substituted with 1-4
15 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine,

wherein substituted C₂₋₄-alkenyl is substituted with 1-2 substituents selected from cyano, amino-carbonyl and C₁₋₄-alkyl-oxy-carbonyl, or

A signifies a group of formula Z-CH₂-Y-CH₂-,

20 wherein Y represents O or NR wherein R is hydrogen or C₁₋₄-alkyl and

Z represents heterocyclyl or optionally substituted aryl with 1-3 benzyl-oxy groups;

X represents S or O;

with the proviso that only one of R¹ and R² is hydrogen.

25

3. Compounds as claimed in claim 1 or 2 wherein

- 73 -

R¹ is C₁₋₁₂-alkyl, aryl or C₁₋₄-alkyl substituted with phenyl optionally substituted with 1-5 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine;

5 R² is C₂₋₄-alkenyl, C₁₋₁₂-alkyl or C₁₋₄-alkyl substituted with phenyl optionally substituted with 1-5 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine;

R³ is C₁₋₁₂-alkyl, C₃₋₈-cycloalkyl, phenyl, benzyl or heterocyclyl,
wherein a phenyl ring may be substituted with 1-5 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine, bromine and
10 cyano;

A is C₁₋₇-alkyl, hydroxy-C₁₋₄-alkyl, C₂₋₄-alkenyl, substituted C₂₋₄-alkenyl, aryl-CH(OH)-, C₁₋₄-alkyl substituted with C₁₋₄-alkoxy, C₁₋₄-alkyl-carbonyl-oxy-, heterocyclyl-oxy or A is C₁₋₄-alkyl substituted with phenyl optionally substituted with 1-5 substituents or with heterocyclyl optionally substituted with 1-4
15 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine,

wherein substituted C₂₋₄-alkenyl is substituted with 1-2 substituents selected from cyano, amino-carbonyl and C₁₋₄-alkyl-oxy-carbonyl, or

A signifies a group of formula Z-CH₂-Y-CH₂-,

20 wherein Y represents O or NR wherein R is hydrogen or C₁₋₄-alkyl and

Z represents heterocyclyl or optionally substituted aryl with 1-3 benzyl-oxy groups;

X represents S or O.

25 4. Compounds as claimed in any one of claims 1 to 3 wherein

R¹ is C₁₋₇-alkyl, phenyl or C₁₋₄-alkyl substituted with phenyl;

- 74 -

R² is C₂₋₄-alkenyl, C₁₋₇-alkyl or C₁₋₄-alkyl substituted with phenyl optionally substituted with 1-5 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine;

R³ is C₁₋₇-alkyl, C₃₋₆-cycloalkyl, phenyl, benzyl or heterocyclyl,

5 wherein a phenyl ring may be substituted with 1-5 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine, bromine and cyano;

A is C₁₋₇-alkyl, hydroxy-C₁₋₄-alkyl, C₂₋₄-alkenyl, substituted C₂₋₄-alkenyl, phenyl-CH(OH)-, C₁₋₄-alkyl substituted with C₁₋₄-alkoxy, C₁₋₄-alkyl-carbonyl-oxy-,
10 heterocyclyl-oxy or A is C₁₋₄-alkyl substituted with phenyl optionally substituted with 1-5 substituents or with pyridyl optionally substituted with 1-4 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine,

 wherein substituted C₂₋₄-alkenyl is substituted with 1-2 substituents selected from cyano, amino-carbonyl and C₁₋₄-alkyl-oxy-carbonyl, or

15 A signifies a group of formula Z-CH₂-Y-CH₂-,

 wherein Y represents O or NR wherein R is hydrogen or C₁₋₄-alkyl and

 Z represents pyridyl or optionally substituted phenyl with 1-3 benzyl-oxy groups;

 X represents S or O.

20

5. Compounds as claimed in any one of claims 1 to 4 wherein

 R¹ is C₁₋₇-alkyl or C₁₋₄-alkyl substituted with phenyl;

 R² is C₂₋₄-alkenyl, C₁₋₇-alkyl or C₁₋₄-alkyl substituted with phenyl;

 R³ is C₁₋₄-alkyl, C₃₋₆-cycloalkyl or optionally substituted phenyl,

25 wherein phenyl may be substituted with 1-5 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine, bromine and cyano;

A is C₁₋₇-alkyl, C₂₋₄-alkenyl, substituted C₂₋₄-alkenyl, C₁₋₄-alkyl substituted with phenyl optionally substituted with 1-5 substituents or with pyridyl optionally substituted with 1-4 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine,

5 wherein substituted C₂₋₄-alkenyl is substituted with 1-2 substituents selected from cyano, amino-carbonyl and C₁₋₄-alkyl-oxy-carbonyl, or

A signifies a group of formula Z-CH₂-Y-CH₂-,

 wherein Y represents O or NR wherein R is hydrogen or C₁₋₄-alkyl and

 Z represents pyridyl or optionally substituted phenyl with 1-3 benzyl-oxy
10 groups;

X represents S or O.

6. Compounds as claimed in any one of claims 1 to 5 wherein

R¹ is C₁₋₇-alkyl;

15 R² is C₁₋₇-alkyl;

R³ is optionally substituted phenyl,

 wherein phenyl may be substituted with 1-5 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine, bromine and cyano;

20 A is C₁₋₄-alkyl substituted with phenyl optionally substituted with 1-3 substituents or with pyridyl optionally substituted with 1-2 substituents selected from C₁₋₄-alkyl, C₁₋₄-alkoxy, hydroxy, fluorine, chlorine and bromine, or

A signifies a group of formula Z-CH₂-Y-CH₂-,

 wherein Y represents O or NR wherein R is hydrogen or C₁₋₄-alkyl and

 Z represents pyridyl or phenyl;

25 X represents S or O.

7. Compounds as claimed in any one of claims 1 to 6 wherein

R^1 is C_{1-4} -alkyl;

R^2 is C_{1-4} -alkyl;

5 R^3 is optionally substituted phenyl,

wherein phenyl may be substituted with 1-3 substituents selected from C_{1-2} -alkyl, C_{1-2} -alkoxy, hydroxy, fluorine, chlorine, bromine and cyano;

A is C_{1-2} -alkyl substituted with phenyl optionally substituted with 1-3 substituents or with pyridyl optionally substituted with 1-2 substituents selected from C_{1-4} -alkyl,
10 C_{1-4} -alkoxy, hydroxy, fluorine, chlorine and bromine, or

A signifies a group of formula $Z-CH_2-Y-CH_2-$,

wherein Y represents O or NR wherein R is hydrogen or C_{1-2} -alkyl and

Z represents pyridyl or phenyl;

X represents S or O.

15

8. Compounds as claimed in any one of claims 1 to 7 wherein

R^1 is isopropyl;

R^2 is methyl or ethyl;

R^3 is optionally substituted phenyl,

20 wherein phenyl may be substituted with 1-3 chlorine substituents;

A is C_{1-2} -alkyl substituted with phenyl or pyridyl, or

A signifies a group of formula $Z-CH_2-Y-CH_2-$,

wherein Y represents O or NR wherein R is hydrogen or C_{1-2} -alkyl and

- 77 -

Z represents pyridyl or phenyl;

X represents S or O.

9. Compounds as claimed in any one of claims 1 to 8 wherein

5 R^1 is isopropyl;

R^2 is methyl;

R^3 is optionally substituted phenyl,

wherein phenyl may be substituted with 1-2 chlorine substituents;

A is C_{1-2} -alkyl substituted with pyridyl, or


10 A signifies a group of formula $Z-CH_2-Y-CH_2-$,

wherein Y represents O or NR wherein R is hydrogen or C_{1-2} -alkyl and

Z represents pyridyl;

X represents S or O.

15 10. Compounds of formula I



wherein

R^1 is hydrogen, C_{1-12} -alkyl, C_{3-8} -cycloalkyl, aryl or C_{1-4} -alkyl substituted with phenyl optionally substituted with 1-5 substituents selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine and bromine;

20

- 78 -

R^2 is hydrogen, C_{1-12} -alkyl or C_{1-4} -alkyl substituted with phenyl optionally substituted with 1-5 substituents selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine and bromine;

R^3 is C_{1-12} -alkyl, C_{3-8} -cycloalkyl, phenyl or benzyl,

5 wherein a phenyl ring may be substituted with 1-5 substituents selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine and bromine;

A is hydroxy- C_{1-4} -alkyl or aryl-CH(OH)-; or

A signifies a group of formula Z-CH₂-Y-CH₂-,

 wherein Y represents O or NR wherein R is hydrogen or C_{1-4} -alkyl and

10 Z represents aryl or heterocyclyl; or

A signifies C_{1-4} -alkyl substituted with phenyl optionally substituted with 1-5 substituents or with heterocyclyl optionally substituted with 1-4 substituents selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine and bromine;

X represents S or O;

15 with the proviso that only one of R^1 and R^2 is hydrogen;

as well as hydrolyzable esters or ethers of compounds of formula I and pharmaceutically acceptable salts thereof.

11. Compounds as claimed in any one of claims 1 to 10 wherein

20 X represents S.

12. Compounds as claimed in claims 1 or 10 which compounds are selected from:

4-(3,5-dichlorophenylthio)-5-hydroxymethyl-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone,


- 5-(benzylaminomethyl)-4-(3,5-dichlorophenylthio)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone,
- 4-(3,5-dichlorophenylthio)-5-(alpha(RS)-hydroxybenzyl)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone,
- 5 4-(3,5-dichlorophenylthio)-5-hydroxymethyl-1-methyl-3-phenyl-1,3-dihydro-2-imidazolone,
- 5-benzyloxymethyl-4-(3,5-dichlorophenylthio)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone,
- 4-(3,5-dichlorophenylthio)-3-isopropyl-1-methyl-5-[[pyridin-4-ylmethyl]amino]methyl}-1,3-dihydro-2-imidazolone,
- 10 5-(benzylamino-methyl)-3-isopropyl-1-methyl-4-phenylthio-1,3-dihydro-2-imidazolone,
- 3-isopropyl-1-methyl-4-phenylthio-5-[[pyridin-4-ylmethyl]amino]methyl}-1,3-dihydro-2-imidazolone,
- 5-benzyloxymethyl-4-(3,5-dichlorophenoxy)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone,
- 15 5-benzyloxymethyl-4-(3,5-dichlorophenylthio)-1-methyl-3-phenyl-1,3-dihydro-2-imidazolone,
- 4-(3,5-dichlorothio)-3-isopropyl-1-methyl-5-(pyridin-4-ylmethoxymethyl)-1,3-dihydro-2-imidazolone,
- 20 5-(benzylaminomethyl)-4-(3,5-dimethylphenylthio)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone,
- 4-cyclohexylthio-3-isopropyl-1-methyl-5-[[pyridin-4-ylmethyl]amino]methyl}-1,3-dihydro-2-imidazolone,
- 4-(3,5-dimethylphenylthio)-3-isopropyl-1-methyl-5-[[pyridin-4-ylmethyl]amino]methyl}-1,3-dihydro-2-imidazolone,
- 25 4-benzylthio-3-isopropyl-1-methyl-5-[[pyridin-4-ylmethyl]-amino]-methyl}-1,3-dihydro-2-imidazolone,

- 4-(3-chlorophenylthio)-3-isopropyl-1-methyl-5-[[pyridin-4-ylmethyl]amino]methyl}-
1,3-dihydro-2-imidazolone,
- 3-isopropyl-4-isopropylthio-1-methyl-5-[[pyridin-4-ylmethyl]amino]methyl}-1,3-
dihydro-2-imidazolone,
- 5 4-isobutylthio-3-isopropyl-1-methyl-5-[[pyridin-4-ylmethyl]amino]methyl}-1,3-
dihydro-2-imidazolone,
- 4-(3,5-dichlorophenylthio)-3-isopropyl-1-(4-methoxybenzyl)-5-[[pyridin-4-
ylmethyl]amino]methyl}-1,3-dihydro-2-imidazolone,
- 4-(3,5-dichlorophenylthio)-5-hydroxymethyl-3-isopropyl-1-(4-methoxybenzyl)-1,3-
10 dihydro-2-imidazolone,
- 4-(3,5-dichlorophenylthio)-3-isopropyl-1-methyl-5-phenethyl-1,3-dihydro-2-
imidazolone,
- 4-(3,5-dichlorophenylthio)-5-(2-hydroxybenzyl)-3-isopropyl-1-methyl-1,3-dihydro-2-
imidazolone,
- 15 3-benzyl-4-(3,5-dichlorophenylthio)-1-methyl-5-[[pyridin-4-ylmethyl]amino]methyl}-
1,3-dihydro-2-imidazolone,
- 1,3-dibenzyl-4-(3,5-dichlorophenylthio)-5-[[pyridin-4-ylmethyl]amino]methyl}-1,3-
dihydro-2-imidazolone,
- 1-benzyl-4-(3,5-dichlorophenylthio)-3-isopropyl-5-[[pyridin-4-
20 ylmethyl]amino]methyl}-1,3-dihydro-2-imidazolone,
- 1-benzyl-4-(3,5-dichlorophenylthio)-3-propyl-5-[[pyridin-4-ylmethyl]amino]methyl}-
1,3-dihydro-2-imidazolone,
- 5-benzyl-4-(3,5-dichlorophenylthio)-3-isopropyl-1-methyl-1,3-dihydro-2-imidazolone,
- 3-benzyl-4-(3,5-dichlorophenylthio)-5-hydroxymethyl-1-methyl-1,3-dihydro-2-
25 imidazolone,
- 4-(3,5-dichlorophenoxy)-3-isopropyl-1-methyl-5-[[pyridin-4-ylmethyl]amino]methyl}-
1,3-dihydro-2-imidazolone,

- 4-sec-butylthio-3-isopropyl-1-methyl-5-[[pyridin-4-ylmethyl]amino]methyl}-1,3-dihydro-2-imidazolone,
- 3-isopropyl-4-(3-methoxyphenylthio)-1-methyl-5-[[pyridin-4-ylmethyl]amino]methyl}-1,3-dihydro-2-imidazolone,
- 5 3-benzyl-4-(3,5-dichlorophenylthio)-5-(1-hydroxyethyl)-1-methyl-1,3-dihydro-2-imidazolone.
13. Compounds as claimed in claim 1 which compounds are selected from:
- 4-(3,5-Dichloro-phenylsulfanyl)-3-isopropyl-5-methoxymethyl-1-methyl-1,3-dihydro-10 imidazol-2-one,
- 4-Hydroxymethyl-3-methyl-1-phenyl-5-(pyridin-4-ylsulfanyl)-1,3-dihydro-imidazol-2-one,
- 1-Benzyl-4-hydroxymethyl-3-methyl-5-(pyridin-4-ylsulfanyl)-1,3-dihydro-imidazol-2-one; hydrochloric acid,
- 15 4-(3,5-Dichloro-phenylsulfanyl)-3-isopropyl-1-methyl-5-(pyridin-3-ylmethoxymethyl)-1,3-dihydro-imidazol-2-one,
- 4-(4-Benzyloxy-benzyloxymethyl)-5-(3,5-dichloro-phenylsulfanyl)-1-isopropyl-3-methyl-1,3-dihydro-imidazol-2-one,
- Acetic acid 5-(3,5-dichloro-phenylsulfanyl)-1-isopropyl-3-methyl-2-oxo-2,3-dihydro-1H-20 imidazol-4-ylmethyl ester,
- 4-(3,5-Dichloro-phenylsulfanyl)-3-isopropyl-1-methyl-5-(pyridin-4-ylloxymethyl)-1,3-dihydro-imidazol-2-one,
- 3-(5-Benzyl-3-isopropyl-1-methyl-2-oxo-2,3-dihydro-1H-imidazol-4-ylsulfanyl)-benzonitrile,
- 25 4-Benzyl-5-isobutylsulfanyl-1-isopropyl-3-methyl-1,3-dihydro-imidazol-2-one,
- 4-Benzyl-1-isopropyl-5-isopropylsulfanyl-3-methyl-1,3-dihydro-imidazol-2-one,
- 4-Benzyl-1-isopropyl-3-methyl-5-methylsulfanyl-1,3-dihydro-imidazol-2-one,

- 4-(3,5-Dichloro-phenylsulfanyl)-5-isobutyl-3-isopropyl-1-methyl-1,3-dihydro-imidazol-2-one,
- 4-Benzyl-5-(3,5-dichloro-phenylsulfanyl)-3-ethyl-1-isopropyl-1,3-dihydro-imidazol-2-one,
- 5 1-Allyl-5-benzyl-4-(3,5-dichloro-phenylsulfanyl)-3-isopropyl-1,3-dihydro-imidazol-2-one,
- 4-(3,5-Dichloro-phenylsulfanyl)-1-ethyl-3-isopropyl-5-pyridin-4-ylmethyl-1,3-dihydro-imidazol-2-one,
- 1-Allyl-4-(3,5-dichloro-phenylsulfanyl)-3-isopropyl-5-pyridin-4-ylmethyl-1,3-dihydro-
- 10 imidazol-2-one,
- [E]-3-[5-(3,5-Dichloro-phenylsulfanyl)-1-isopropyl-3-methyl-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-acrylic acid ethyl ester,
- 3-[5-(3,5-Dichloro-phenylsulfanyl)-1-isopropyl-3-methyl-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-acrylonitrile,
- 15 3-[5-(3,5-Dichloro-phenylsulfanyl)-1-isopropyl-3-methyl-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-acrylamide,
- 1-Benzyl-5-(3,5-dichloro-phenylsulfanyl)-3-methyl-4-vinyl-1,3-dihydro-imidazol-2-one, or
- 4-(3,5-Dichloro-phenylsulfanyl)-3-isopropyl-1-methyl-5-pyridin-4-ylmethyl-1,3-dihydro-
- 20 imidazol-2-one.


14. A process for the preparation of compounds of formula



wherein R¹, R², R³ and X are as defined in formula I;

comprises

reacting chloro compound of formula VII




wherein R^1 and R^2 are as defined in formula I

5 with nucleophilic agent R^3XH

wherein R^3 and X are as defined in formula I.

15. Compounds of formula VIII-a



10 wherein R^1 , R^2 and X are as defined in formula I and R^3 is C_{7-12} -alkyl, C_{3-8} -cycloalkyl, optionally substituted phenyl or optionally substituted benzyl

wherein phenyl or benzyl may be substituted with 1-5 substituents selected from C_{1-4} -alkyl, C_{1-4} -alkoxy, hydroxy, fluorine, chlorine or bromine.

15 16. A compound of formula I, or hydrolyzable ester, ether or pharmaceutically acceptable salt thereof, or a composition containing a compound of formula I, as claimed in any one of claims 1 to 13 for use as medicament.

17. Use of a compound of formula I, or hydrolyzable ester, ether or pharmaceutically acceptable salt thereof, or a composition containing a compound of formula I, as defined in any one of claims 1 to 13 for the preparation of a medicament for the treatment of diseases mediated by the human immunodeficiency virus (HIV).

5

18. A compound of formula I, or hydrolyzable ester, ether or pharmaceutically acceptable salt thereof, or a composition containing a compound of formula I as claimed in any one of claims 1 to 13 for its use in the treatment of a disease mediated by the human immunodeficiency virus (HIV).

10

19. A pharmaceutical composition comprising a pharmaceutically effective amount of a compound of formula I, or hydrolyzable ester, ether or pharmaceutically acceptable salt thereof, as defined in any one of claims 1 to 13 and, if desired, a pharmaceutical inert carrier.

15

20. A pharmaceutical composition according to claim 19 for its use in the treatment of diseases mediated by the human immunodeficiency virus (HIV).

21. The invention as hereinbefore described.

20

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/13069

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D233/86 C07D233/78 C07D401/12 C07D401/06 A61K31/4164
A61K31/4174 A61K31/4427

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, BEILSTEIN Data, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 94 10168 A (MERCK SHARP & DOHME ; MACLEOD ANGUS MURRAY (GB)) 11 May 1994 (1994-05-11) cited in the application the whole document	1-22
X	REISCH, JOHANNES: "Photochemical transformation of pyramidon in the presence of primary alcohols" CHIM. THER. (1966), (5-6), 335-6 , XP001055699 see compounds 3a,3b	1-4,10
Y	EP 0 700 911 A (SHIONOGI & CO) 13 March 1996 (1996-03-13) see whole document, especially definitions in claim 1, and the Synthetic routes, pages 4-6	1-22

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
 E earlier document but published on or after the international filing date
 L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 O document referring to an oral disclosure, use, exhibition or other means
 P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* & * document member of the same patent family

Date of the actual completion of the international search

31 January 2002

Date of mailing of the international search report

20/02/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Scruton-Evans, I

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/13069

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 552 060 A (SHIONOGI & CO) 21 July 1993 (1993-07-21) see whole document, especially definitions in claim 1 and the synthetic routes -----	1-22

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/13069

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9410168	A	11-05-1994	AU 5283593 A	24-05-1994
			WO 9410168 A1	11-05-1994
			US 5612362 A	18-03-1997
EP 0700911	A	13-03-1996	EP 0700911 A1	13-03-1996
			AT 173470 T	15-12-1998
			AT 200082 T	15-04-2001
			AU 649906 B2	02-06-1994
			AU 3182793 A	22-07-1993
			CA 2087112 A1	17-07-1993
			DE 69322106 D1	24-12-1998
			DE 69322106 T2	29-04-1999
			DE 69330072 D1	03-05-2001
			DE 69330072 T2	23-08-2001
			DK 552060 T3	02-08-1999
			DK 700911 T3	07-05-2001
			EP 0552060 A1	21-07-1993
			ES 2125301 T3	01-03-1999
			ES 2157291 T3	16-08-2001
			HU 63615 A2	28-09-1993
			JP 5255270 A	05-10-1993
			KR 259309 B1	01-08-2000
			NZ 245674 A	27-06-1995
			PT 700911 T	31-07-2001
			US 5326780 A	05-07-1994
			US 5472965 A	05-12-1995
			ZA 9300217 A	18-08-1993
EP 0552060	A	21-07-1993	AT 173470 T	15-12-1998
			AT 200082 T	15-04-2001
			AU 649906 B2	02-06-1994
			AU 3182793 A	22-07-1993
			CA 2087112 A1	17-07-1993
			DE 69322106 D1	24-12-1998
			DE 69322106 T2	29-04-1999
			DE 69330072 D1	03-05-2001
			DE 69330072 T2	23-08-2001
			DK 552060 T3	02-08-1999
			DK 700911 T3	07-05-2001
			EP 0552060 A1	21-07-1993
			EP 0700911 A1	13-03-1996
			ES 2125301 T3	01-03-1999
			ES 2157291 T3	16-08-2001
			HU 63615 A2	28-09-1993
			JP 5255270 A	05-10-1993
			KR 259309 B1	01-08-2000
			NZ 245674 A	27-06-1995
			PT 700911 T	31-07-2001
			US 5326780 A	05-07-1994
			US 5472965 A	05-12-1995
			ZA 9300217 A	18-08-1993